

Asymptotic Dynamics of Nonlinear Schrödinger Equations with Many Bound States

TAI-PENG TSAI

Institute for Advanced Study, Princeton, NJ 08540

E-mail: ttsai@ias.edu

Abstract

We consider a nonlinear Schrödinger equation with a bounded local potential in $\mathbb{R}^3$. The linear Hamiltonian is assumed to have three or more bound states with the eigenvalues satisfying some resonance conditions. Suppose that the initial data is localized and small of order n in H^1 , and that its ground state component is larger than $n^{3-\epsilon}$ with $\epsilon > 0$ small. We prove that the solution will converge locally to a nonlinear ground state as the time tends to infinity.

1 Introduction

Consider the nonlinear Schrödinger equation

$$i\partial_t\psi = (-\Delta + V)\psi + \lambda|\psi|^2\psi, \quad \psi(t=0) = \psi_0, \quad (1.1)$$

where $V = V(x)$ is a smooth localized real potential, $\lambda = \pm 1$ and $\psi = \psi(t, x) : \mathbb{R} \times \mathbb{R}^3 \longrightarrow \mathbb{C}$ is a wave function. For any solution $\psi(t) \in H^1(\mathbb{R}^3)$ the L^2 -norm and the Hamiltonian

$$\mathcal{H}[\psi] = \int \frac{1}{2}|\nabla\psi|^2 + \frac{1}{2}V|\psi|^2 + \frac{1}{4}\lambda|\psi|^4 dx \quad (1.2)$$

are constant for all t . The global well-posedness for small solutions in $H^1(\mathbb{R}^3)$ can be proven using these conserved quantities and a continuity argument, no matter what the sign of λ is. We assume that the linear Hamiltonian $H_0 := -\Delta + V$ has $N+1$ simple eigenvalues $e_0 < e_1 < \dots < e_N$ with normalized eigenvectors ϕ_k , $k = 0, 1, \dots, N$, where $N \geq 2$. These eigenvalues are assumed to satisfy some resonance conditions to be specified later on. The *nonlinear bound states* to the Schrödinger equation (1.1) are solutions to the equation

$$(-\Delta + V)Q + \lambda|Q|^2Q = EQ, \quad (1.3)$$

¹2000 *Mathematics Subject Classification*. Primary. 35Q40, 35Q55.

²*Key words and phrases*. asymptotic dynamics, Schrödinger equations, many bound states.

for some E . They are critical points to the Hamiltonian $\mathcal{H}[\psi]$ defined in (1.2) subject to the constraint of fixed L^2 -norm. For any nonlinear bound state $Q = Q_E(x)$, the function $\psi(t, x) = Q(x)e^{-iEt}$ is an exact solution to the nonlinear Schrödinger equation (1.1). We may obtain $N + 1$ families of such nonlinear bound states by standard bifurcation theory, corresponding to the $N + 1$ eigenvalues of the linear Hamiltonian. For any E sufficiently close to e_0 so that $E - e_0$ and λ have the same sign, there is a unique positive solution $Q = Q_E$ to (1.3) which decays exponentially as $x \rightarrow \infty$. We call this family the *nonlinear ground states* and we refer to it as $\{Q_E\}_E$. Similarly, for each $k > 0$ there is a *nonlinear excited state* family $\{Q_{k,E_k}\}_{E_k}$ for E_k near e_k . These solutions are small and $\|Q_{k,E_k}\| \sim |E_k - e_k|^{1/2}$. See Lemma 2.1.

Our goal is to understand the long-time dynamics of the solutions at the presence of nonlinear bound states. One first considers the stability of nonlinear ground states. There are two different concepts: *orbital stability* and *asymptotic stability*. It is well-known that nonlinear ground states are orbitally stable in the sense that the difference

$$\inf_{\Theta, E} \|\psi(t) - Q_E e^{i\Theta}\|_{L^2(\mathbb{R}^3)}$$

remains small for all time t if it is initially small. On the other hand, one expects that the difference actually approaches zero locally, as the majority of the difference is a dispersive wave which escapes to infinity. Hence one expects that it is asymptotically stable in the sense that

$$\|\psi(t) - Q_{E(t)} e^{i\Theta(t)}\|_{L^2_{\text{loc}}} \longrightarrow 0$$

as $t \rightarrow \infty$, for a suitable choice of $E(t)$ and $\Theta(t)$. Here $\|\cdot\|_{L^2_{\text{loc}}}$ denotes a local L^2 norm, a precise choice will be made later on in (1.8). One also wants to determine the decay rate and whether $E(t)$ has a limit. A more difficult problem, which one studies next, is the asymptotic dynamics of the solutions when the initial data are away from nonlinear ground states. As in the previous problem, one wants to identify their local behavior.

If $-\Delta + V$ has only one bound state, i.e., with no excited states, the asymptotic stability is proved in [16]. The solution eventually settles down to some ground state Q_{E_∞} with E_∞ close to E and the local difference is bounded by $Ct^{-3/2}$, as the decay rate of the free evolution $e^{it\Delta}$. This result is extended in [10] to all small initial data, not necessarily near ground states. See also [1, 4].

Suppose $-\Delta + V$ has two bound states. We proved in [19] that the evolution with initial data ψ_0 near some Q_E will eventually settle down to some ground state Q_{E_∞} with E_∞ close to E . The local difference is, however, only bounded by $Ct^{-1/2}$ due to the persistence of the excited state. The key mechanism here is the resonance decay of the excited state due to resonance with the continuous spectrum. See [2, 3]

for an one dimensional equation, [5] its extension to higher dimensions, and [17] for real-valued nonlinear Klein-Gordon equations.

The problem becomes more delicate when the initial data are away from ground states. Based on physical intuition, one expects that all solutions near excited states decay to ground states unless initially they are exactly excited states. However, we proved [21] that there exists a family of “finite co-dimensional manifolds” in the space of initial data so that the dynamics asymptotically converge to some excited states. Outside a small neighborhood of these manifolds, the asymptotic profiles are given by some ground states [20]. We further showed [22] that there are exactly three asymptotic profiles: vacuum, excited states or ground states. The last problem is also considered in [18]. Earlier works concerning related linear analysis were obtained in [6, 7, 13, 14, 23].

In this paper, we extend the results in [19, 20] to the case when $-\Delta + V$ has three or more bound states.

Our assumptions on the operator $-\Delta + V$ are as follows:

Assumption A0: $-\Delta + V$ acting on $L^2(\mathbb{R}^3)$ has $N + 1$ simple eigenvalues $e_0 < e_1 < \dots < e_N < 0$, $N \geq 2$, with normalized eigenvectors $\phi_0, \dots, \phi_N$.

Assumption A1: $V(x)$ is a real-valued function. For λQ_E^2 sufficiently small, the bottom of the continuous spectrum to $-\Delta + V + \lambda Q_E^2$, 0, is not a generalized eigenvalue, i.e., not an eigenvalue nor a resonance. Also, we assume that V satisfies the assumption in [24] so that the $W^{k,p}$ estimates $k \leq 2$ for the wave operator $W_{H_0} = \lim_{t \rightarrow \infty} e^{itH_0} e^{it\Delta}$ hold for $k \leq 2$, i.e., there is a small $s_1 > 0$ such that,

$$|\nabla^\alpha V(x)| \leq C \langle x \rangle^{-5-s_1}, \quad \text{for } |\alpha| \leq 2.$$

Also, the functions $(x \cdot \nabla)^k V$, for $k = 0, 1, 2, 3$, are $-\Delta$ bounded with a $-\Delta$ bound less than 1:

$$\|(x \cdot \nabla)^k V \phi\|_2 \leq s_2 \|-\Delta \phi\|_2 + C \|\phi\|_2, \quad s_2 < 1, \quad k = 0, 1, 2, 3.$$

Assumption A2: Resonance condition. Denote $e_{k0} = e_k - e_0$. We assume that

$$e_0 < 2e_1. \tag{1.4}$$

Hence $2e_{k0} > |e_0|$ for all $k > 0$. We further assume that, for some small $s_0 > 0$,

$$\gamma_0 \equiv \inf_{\substack{1 \leq k \leq N \\ |s| < s_0}} \lim_{r \rightarrow 0+} \text{Im} \left(\phi_0 \phi_k^2, \frac{1}{-\Delta + V + e_0 - 2e_k - s - ri} \mathbf{P}_c \phi_0 \phi_k^2 \right) > 0. \tag{1.5}$$

Assumption A3: No-resonance condition. Let $j_{\max} = 3$. For all $j = 2, \dots, j_{\max}$ and for all $k_1, \dots, k_j, l_1, \dots, l_j \in \{0, \dots, N\}$ with $\{k_1, \dots, k_j\} \neq \{l_1, \dots, l_j\}$ as sets with multiplicities, (e.g., $\{0, 0, 1\} \neq \{0, 1, 1\}$),

$$e_{k_1} + \dots + e_{k_j} \neq e_{l_1} + \dots + e_{l_j}. \tag{1.6}$$

Assumption A1 contains some standard conditions to assure that most tools for linear Schrödinger operators apply. These conditions are certainly not optimal. (For example, it is sufficient to assume 0 is not a resonance or eigenvalue of H_0 , which implies the same statement for $-\Delta + V + \lambda Q^2$. See [21].) The main assumption in A2 is the condition $e_0 < 2e_1$. Since the expression for γ_0 is quadratic, it is non-negative and $\gamma_0 > 0$ holds generically. The condition $e_0 < 2e_1$ states that the energies of the excited states are closer to the continuum spectrum than to the ground state energy. It guarantees that, for each $k > 0$, $2e_k - e_0 > 0$ becomes a resonance in the continuum spectrum of H_0 . ($H_0 + e_0 - 2e_k$ is not invertible in L^2 .) This resonance produces the main relaxation/growth mechanism. If this condition fails, the resonance occurs in higher order terms and a proof of relaxation will be much more complicated. Also, the rate of decay will be different. Assumption A3 is a new condition to avoid direct resonance between the eigenvalues. It is trivial if $N = 0, 1$. It holds true generically and is often seen in dynamical systems of ODE's. See Example 2 at the end of this section for what may happen if this assumption fails. If we relax the assumption $e_0 < 2e_1$, we may need to increase $j_{\max}$.

Denote by L_r^2 the weighted L^2 spaces (r may be positive or negative),

$$L_r^2(\mathbb{R}^3) = \{\phi \in L^2(\mathbb{R}^3) : (1 + x^2)^{r/2} \phi \in L^2(\mathbb{R}^3)\}. \quad (1.7)$$

Fix $r_1 > 3$ large enough, to be determined by estimates (2.5) and (2.30). We denote by L_{loc}^p , $p = 1, 2$, the local L^p spaces given by the norm

$$\|\phi\|_{L_{\text{loc}}^p(\mathbb{R}^3)} \equiv \left\{ \int_{\mathbb{R}^3} (1 + |x|)^{-2r_1} |\phi|^p d^3x \right\}^{1/p}. \quad (1.8)$$

Now we are ready to state our main theorem.

Theorem 1.1 *Suppose the assumptions A0–A3 on $H_0 = -\Delta + V$ hold and let $\epsilon > 0$ be any small constant. There is a small constant $n_0 > 0$ such that the following holds. Let $\psi(t, x)$ be a solution of (1.1) with initial data ψ_0 decomposed as*

$$\psi_0 = x_0^0 \phi_0 + x_1^0 \phi_1 + \cdots + x_N^0 \phi_N + \xi_0$$

with respect to H_0 , where $x_k^0 \in \mathbb{C}$, $k = 0, 1, \dots, N$, and $\xi_0 \in \mathbf{H}_c(H_0)$. Suppose that

$$\|\psi_0\|_Y = n, \quad 0 < n \leq n_0, \quad |x_0^0| \geq n^{3-\epsilon}, \quad \|\xi_0\|_Y \leq n/2, \quad (1.9)$$

where Y is the space for initial data,

$$Y \equiv H^1 \cap L^1(\mathbb{R}^3). \quad (1.10)$$

Then there exists an E_∞ with $\|Q_{E_\infty}\|_{L^2} \sim n$ and a real function $\Theta(t) = -E_\infty t + O(\log t)$ such that

$$\|\psi(t) - Q_{E_\infty} e^{i\Theta(t)}\|_{L^2_{\text{loc}}} \leq C_2(1+t)^{-1/2}, \quad (1.11)$$

for some constant $C_2 > 0$ depending on n . Suppose, furthermore, $\max_{k=1}^N |x_k^0| \geq n/100$, we also have a lower bound

$$\|\psi(t) - Q_{E_\infty} e^{i\Theta(t)}\|_{L^2_{\text{loc}}} \geq C_1(1+t)^{-1/2}, \quad (1.12)$$

for some constant $C_1 > 0$ depending on n .

The condition $|x_0^0| \geq n^{3-\epsilon}$ is certainly not optimal and can be greatly relaxed. It ensures that ψ_0 is away from nonlinear excited states and that the dispersion component, ξ_0 , is not extremely large compared with x_0^0 . This condition, however, still allows the ground state component to be much smaller than other components and exhibit the main phenomena.

Depending on the relative sizes of the bound states, there are three regimes:

- I. when an excited state is dominant;
- II. when the ground state and the excited states are comparable;
- III. when the ground state is dominant.

Because the dominant terms are different in different regimes, the natural linear operators and the corresponding decompositions of the wave function are different. In regime II we can use H_0 as the linear operator and decompose a function $\psi \in L^2$ as

$$\psi = x_0\phi_0 + \cdots + x_N\phi_N + \xi, \quad (1.13)$$

with $x_k \in \mathbb{C}$ and $\xi \in \mathbf{H}_c(H_0)$. When the function ψ is close to a nonlinear ground state Q_E , i.e., in regime III, it is natural to use $\mathcal{L} = \mathcal{L}_E$, the linearized operator around Q_E ,

$$\mathcal{L}h = -i \{ (-\Delta + V - E + 2\lambda Q^2) h + \lambda Q^2 \bar{h} \}, \quad (1.14)$$

and decompose the wave function according to the spectral decomposition with respect to $\mathcal{L}$. Notice that $\mathcal{L}$ is not self-adjoint. Similarly, we can use linearized operators around excited states in regime I.

The picture of the dynamics is as follows. Suppose the ground state component is initially of order $n^{3-\epsilon}$ and the solution is in regime II. The ground state component

will gradually grow while the excited states gradually decay, until the solution enters regime III, i.e., when the ground state component becomes much larger than other components. This time interval is called the *transition regime*. After entering regime III, the solution will converge locally to some nonlinear ground state as time tends to infinity, with the excited state components vanishing and the dispersion component escaping to infinity. This time interval is called the *stabilization regime*.

As indicated above, we will use different operators and coordinate systems for these two regimes. Besides technical problems associated with changing coordinate systems, there is an intrinsic difficulty related to the time reversibility of the Schrödinger equations; It is not sufficient to control only the usual Sobolev space norms at the time of coordinates changing. To overcome this difficulty, we will use a concept introduced in [20, 22], the *out-going estimates*, to capture the time-direction sensitive information of the dispersive waves.

As it will be seen from examples below, there are energy transfers from higher modes to lower modes. Although all excited states eventually vanish, in an intermediate time an excited state may actually grow because it gains more energy than it loses. This complicates the analysis: one cannot prove the decay of each individual excited state for all time. Instead, we have derived some *monotonicity formulas* for their sum, see (4.57). Because the energy transfer between excited states is relatively small in the stabilization regime, the phenomenon mentioned above is only apparent in the transition regime.

We now give two examples illustrating some phenomena of many bound states, in particular the relaxation/growth mechanism. We first recall the concept of *phase factor*. In our analysis it is essential to identify the main oscillation factors of various terms. If we decompose the solution $\psi(t)$ according to (1.13), the equation for a component $x_k(t)$ is

$$i\dot{x}_k = e_k x_k + (\phi_k, \lambda \psi^2 \bar{\psi}). \quad (1.15)$$

From the linear part we find that $x_k(t)$ has an oscillation factor $e^{-ie_k t}$. We write $x_k(t) = e^{-ie_k t} u_k(t)$ and say that its phase factor is $-e_k$. We will talk about phase factors of polynomials in x_k in a similar way. For example, the phase factor of $x_l x_m \bar{x}_j$ is $-e_l - e_m + e_j$.

Example 1. Three bound states case in transition regime

Suppose H_0 has three bound states and denote $x(t) = x_0(t)$, $y(t) = x_1(t)$ and $z(t) = x_2(t)$. The leading terms of ξ are generated by bound states and are cubic in x_j . Those with resonant coefficient functions, i.e., those with *negative phase factors*,

give the relevant part of ξ ,

$$\xi = (y^2 + z^2 + yz)\bar{x} + z^2\bar{y} + \dots \quad (1.16)$$

Here we have ignored their coefficient functions, which are non-local and complex because of the resonance with the continuous spectrum. Denote $\chi = \sum_{j=0}^N x_j \phi_j = x\phi_0 + y\phi_1 + z\phi_2$. The main terms of (1.15) for x_k are

$$i\dot{x}_k = e_k x_k + (\phi_k, \lambda\chi^2\bar{\xi} + 2\lambda\chi\bar{\chi}\xi) + \dots, \quad (1.17)$$

where $\dots$ denotes irrelevant terms. Substituting (1.16) into (1.17), we get

$$\begin{aligned} \dot{x} &= [\gamma_{22}^0 |z|^4 + 4\gamma_{12}^0 |y|^2 |z|^2 + \gamma_{11}^0 |y|^4] x + \dots \\ \dot{y} &= [\gamma_{22}^1 |z|^4 - 4\gamma_{12}^0 |x|^2 |z|^2 - 2\gamma_{11}^0 |x|^2 |y|^2] y + \dots \\ \dot{z} &= [-2\gamma_{22}^0 |x|^2 |z|^2 - 4\gamma_{12}^0 |x|^2 |y|^2 - 2\gamma_{22}^1 |y|^2 |z|^2] z + \dots \end{aligned} \quad (1.18)$$

Here $\dots$ denotes irrelevant and error terms; γ_{11}^0 , γ_{12}^0 , γ_{22}^0 and γ_{22}^1 are non-negative constants to be defined later in (4.26). These constants (except γ_{22}^1 , depending on whether $e_1 - 2e_2 < 0$) are generically positive. They are the nonlinear analogue of the *Fermi golden rule*, extensively studied in, e.g., [15, 17, 2, 3, 5, 21]. The irrelevant terms have two kinds. The first kind consists of terms with different phase factors. They have few effect averaging over time and can be removed using a *normal form* procedure, see Lemma 4.2. The second kind consists of terms with same phase factors but with purely imaginary coefficients. They only contribute to the phase of x_k , not to the magnitude. To illustrate further, let us assume $\gamma_{11}^0 = \gamma_{12}^0 = \gamma_{22}^0 = \gamma_{22}^1 = 1$ and denote $A = |x|^2$, $B = |y|^2$, $C = |z|^2$. By (1.18) we have

$$\begin{aligned} \dot{A}/2 &= 4ABC + AB^2 + AC^2 + \dots \\ \dot{B}/2 &= -4ABC - 2AB^2 + BC^2 + \dots \\ \dot{C}/2 &= -4ABC - 2AC^2 - 2BC^2 + \dots \end{aligned} \quad (1.19)$$

where $\dots$ denotes irrelevant terms. Although the above system is accurate only in the transition regime, corresponding equations for other regimes are similar. From (1.19) we can read the energy transfers from higher modes to lower modes. Only half of the energy decrease in a higher mode goes to lower modes, while the other half goes to dispersion (radiation). The lowest mode (ground state) is only receiving energy while the highest mode (x_N) is only losing energy. The intermediate modes receive energy from higher modes and release energy to lower modes and radiation.

Richer phenomena occur when the constants γ_{lm}^j have different sizes. For example, suppose γ_{22}^1 is much larger than other constants and initially the energy is concentrated in the third mode z . Then the second mode y will first grow exponentially,

acquiring energy from z , and then gradually decay. Another phenomenon to be noticed is the interaction between three modes, the ABC terms in (1.19). They have the same coefficients $\pm 4\gamma_{12}^0$ in (1.18). Hence this interaction is more apparent when γ_{12}^0 is much larger than other coefficients. This phenomenon is present only for many bound states case, $N \geq 2$.

Example 2. No-resonance assumption A3 violated

Still assume $e_0 < 2e_1$ and three bound states. The only possibility for assumption A3 to fail is

$$e_0 + 2e_2 = 3e_1. \quad (1.20)$$

An example is $e_0 = -10$, $e_1 = -4$ and $e_2 = -1$. Note that

$$e_1 - 2e_2 = e_0 - 2e_1 < 0. \quad (1.21)$$

Hence γ_{22}^1 is generically positive. Because of (1.20), when we substitute (1.16) into (1.17) we get new resonant terms in (1.18):

$$\begin{aligned} \dot{x} &= [\dots]x + y^2 \overline{(z^2 \bar{y})} + \dots \\ \dot{y} &= [\dots]y + z^2 \overline{(y^2 \bar{x})} + x\bar{y}(\bar{y}z^2) + \dots \\ \dot{z} &= [\dots]z + y\bar{z}(\bar{x}y^2) + \dots \end{aligned} \quad (1.22)$$

The first group of terms on the right side are those in (1.18). The cubic terms in the parentheses () are from (1.16). For example, $y^2 \overline{(z^2 \bar{y})}$ has the phase factor $-2e_1 - (-2e_2 + e_1) = -e_0$, the same as x , due to (1.20). Although these new terms on the right side have the same phase factors as the left side, their phases are not exactly the same. Moreover, their coefficients are not quadratic and it is unclear how to determine their signs. Hence it is difficult to predict the dynamics of this system.

2 Preliminaries

We first fix the notation. Let H^k denote the Sobolev spaces $W^{k,2}(\mathbb{R}^3)$. The weighted Sobolev space $L_r^2(\mathbb{R}^3)$ is defined in (1.7). Denote by **conj** the conjugation operator. The L^2 inner product $(\cdot, \cdot)$ is

$$(f, g) = \int_{\mathbb{R}^3} \bar{f} g \, d^3x. \quad (2.1)$$

For a function $\phi \in L^2$, denote by $\phi^\perp$ the L^2 -subspace $\{g \in L^2 : (\phi, g) = 0\}$.

In what follows we collect some facts about nonlinear bound states and linear analysis. Their proofs can be found in [19, 20, 21]. Although the proofs there are for two bound states case, the proofs for the many bound states case are the same.

2.1 Nonlinear bound states

Recall that nonlinear bound states of the equation (1.1) are solutions of (1.3). They are critical points of the energy functional $\mathcal{H}[\psi]$ defined in (1.2), subject to the constraint of fixed L^2 -norm. For each such solution Q_E , the function $\psi(t, x) = Q_E(x)e^{-iEt}$ is an exact solution of (1.1). Since we have $N + 1$ simple eigenvalues, we have $N + 1$ families of corresponding nonlinear bound states. The existence and basic properties of these nonlinear bound states are summarized in the following lemma. They are proven in [19, 20] using a contraction mapping argument.

Lemma 2.1 *Suppose $-\Delta + V$ satisfies assumptions A0–A1. Let n_0 be sufficiently small. For each eigenvalue e_k with normalized bound state ϕ_k , $k = 0, 1, \dots, N$, there is a family of nonlinear bound states $\{Q_{k,E_k}\}_{E_k}$ to (1.1) for E_k between e_k and $e_k + \lambda n_0^2$ such that Q_{k,E_k} are real, localized, smooth, and $\lambda^{-1}(E_k - e_k) > 0$,*

$$Q_{k,E_k} = n \phi_k + h, \quad h \perp \phi_k, \quad h = O(n^3) \text{ in } H^2,$$

where $n = [(E_k - e_k)/(\lambda \int \phi_k^4 dx)]^{1/2}$. Moreover, we have $\partial_{E_k} Q_{k,E_k} = O(n^{-2}) Q_{k,E_k} + O(n) = O(n^{-1})$, and $\partial_E^2 Q_{k,E} = O(n^{-3})$. For ground states we will drop the subscript and write Q_E and $R_E = \partial_E Q_E$. We have $R_E = Cn^{-2}Q_E + O(n)$. If we define $c_1 \equiv (Q_E, R_E)^{-1}$, we have $c_1 = O(1)$ and $\lambda c_1 > 0$.

The following lemma summarizes the renormalization results near ground states. They are proven in [19] using implicit function theorem. The L^2 -subspace $M = M_E$ will be defined in Lemma 2.4 (8).

Lemma 2.2 *Let $Y_1 = Y$, defined in (1.10), or $Y_1 = L_{\text{loc}}^2$, defined in (1.8). There are small constants $n_0 > 0$ and $\varepsilon_0 > 0$ such that the following hold. Suppose $\psi \in Y_1$ is close to a nonlinear ground state $Q_E e^{i\Theta}$ with $\|\psi\|_{Y_1} = n$, $\|\psi - Q_E e^{i\Theta}\|_{Y_1} \leq \tau n$, $0 < n \leq n_0$, $0 < \tau \leq \varepsilon_0$.*

(1) *There are unique small $a, \theta \in \mathbb{R}$ and $h \in M_E$ such that*

$$\psi = [Q_E + aR_E + h] e^{i(\Theta + \theta)}. \quad (2.2)$$

Moreover, $\|Q_E\|_{Y_1} \sim n$, $a = O(\tau n^2)$, $h = O(\tau n)$ and $\theta = O(\tau)$.

(2) **(best approximation)** *There is a unique E_* near E such that the component along the R_{E_*} direction as defined by (2.2) vanishes, i.e., there are unique small $\theta_* \in \mathbb{R}$ and $h_* \in M_{E_*}$ such that*

$$\psi = [Q_{E_*} + h_*] e^{i(\Theta + \theta_*)}.$$

Moreover, $E - E_* = O(\tau n^2)$, $h_* = O(\tau n)$ and $\theta_* = O(\tau)$.

(3) Suppose $E' = E + \gamma$ with $|\gamma| \leq \tau^2 n^2$. By part (1) we can rewrite ψ uniquely with respect to E' as

$$\psi = [Q_{E'} + a' R_{E'} + h'] e^{i(\Theta + \theta')},$$

where $h' \in M_{E'}$; h' , a' and θ' are small. We have the estimates

$$E + a - E' - a' = O(\tau\gamma), \quad h - h' = O(n^{-1}\tau\gamma), \quad \theta - \theta' = O(n^{-2}\tau\gamma). \quad (2.3)$$

Notice that $n^{-2}\tau\gamma \leq C\varepsilon_0^3$ is small.

2.2 Linear analysis

We first recall some local decay estimates for e^{-itH_0} . The decay estimate (2.4) is proved in [9, 24] using estimates in [8, 11]. The estimate (2.5) is taken from [17, 19]. The estimate (2.5) holds only if we take $r \rightarrow 0+$, not $r \rightarrow 0-$.

Lemma 2.3 (decay estimates for e^{-itH_0}) Suppose that $H_0 = -\Delta + V$ satisfies the Assumptions A0–A2. For $q \in [2, \infty]$ and $q' = q/(q-1)$,

$$\|e^{-itH_0} \mathbf{P}_c \phi\|_{L^q} \leq C |t|^{-3(\frac{1}{2} - \frac{1}{q})} \|\phi\|_{L^{q'}}. \quad (2.4)$$

For sufficiently large r_1 , for all $k, l, m \in \{0, \dots, N\}$, we have

$$\lim_{r \rightarrow 0+} \left\| \langle x \rangle^{-r_1} \frac{e^{-itH_0}}{(H_0 + e_k - e_l - e_m - ri)} \mathbf{P}_c \langle x \rangle^{-r_1} \phi \right\|_{L^2} \leq C \langle t \rangle^{-3/2} \|\phi\|_{L^2}. \quad (2.5)$$

We now consider the linearized operators around nonlinear ground states. Let $Q = Q_E$ be a nonlinear ground state with $\|Q_E\|_{L^2}$ small. If we consider solutions $\psi(t, x)$ of (1.1) of the form

$$\psi(t, x) = [Q_E(x) + h(t, x)] e^{-iEt},$$

with $h(t, x)$ small in a suitable sense, then $h(t, x)$ satisfies

$$\partial_t h = \mathcal{L}h + \text{nonlinear terms},$$

where the linearized operator $\mathcal{L} = \mathcal{L}_E$ is defined by

$$\mathcal{L}h = -i \{ (-\Delta + V - E + 2\lambda Q^2) h + \lambda Q^2 \bar{h} \}. \quad (2.6)$$

The properties of $\mathcal{L}$ are best understood in the complexification of $L^2(\mathbb{R}^3, \mathbb{C})$.

Definition 2.1 Identify $\mathbb{C}$ with $\mathbb{R}^2$ and $L^2 = L^2(\mathbb{R}^3, \mathbb{C})$ with $L^2(\mathbb{R}^3, \mathbb{R}^2)$. Denote by $\mathbb{C}L^2 = L^2(\mathbb{R}^3, \mathbb{C}^2)$ the complexification of $L^2(\mathbb{R}^3, \mathbb{R}^2)$. $\mathbb{C}L^2$ consists of 2-dimensional vectors whose components are in L^2 . We have the natural embedding

$$\mathbf{j} : f \in L^2 \longrightarrow \begin{bmatrix} \operatorname{Re} f \\ \operatorname{Im} f \end{bmatrix} \in \mathbb{C}L^2.$$

We equip $\mathbb{C}L^2$ with the natural inner product: For $f, g \in \mathbb{C}L^2$, $f = \begin{bmatrix} f_1 \\ f_2 \end{bmatrix}$, $g = \begin{bmatrix} g_1 \\ g_2 \end{bmatrix}$, we define

$$((f, g)) = \int_{\mathbb{R}^3} \bar{f} \cdot g \, d^3x = \int_{\mathbb{R}^3} (\bar{f}_1 g_1 + \bar{f}_2 g_2) \, d^3x. \quad (2.7)$$

Denote by $\mathbf{RE}$ the operator first taking the real part of functions in $\mathbb{C}L^2$ and then pulling back to L^2 :

$$\mathbf{RE} : \mathbb{C}L^2 \rightarrow L^2, \quad \mathbf{RE} \begin{bmatrix} f \\ g \end{bmatrix} = (\operatorname{Re} f) + i(\operatorname{Re} g).$$

We have $\mathbf{RE} \circ \mathbf{j} = \operatorname{id}_{L^2}$.

The operator $\mathcal{L}$ can be naturally extended to an operator acting on $\mathbb{C}L^2$ with the following matrix form:

$$\begin{bmatrix} 0 & L_- \\ -L_+ & 0 \end{bmatrix}, \quad \text{where} \quad \begin{cases} L_- = -\Delta + V - E + \lambda Q^2 \\ L_+ = -\Delta + V - E + 3\lambda Q^2. \end{cases} \quad (2.8)$$

Recall the Pauli matrices

$$\sigma_1 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad \sigma_2 = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}, \quad \sigma_3 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}.$$

They are self-adjoint. We have $\mathbf{RE} \mathcal{L} = \mathcal{L} \mathbf{RE}$ and

$$\sigma_1 \mathcal{L} = \mathcal{L}^* \sigma_1, \quad \sigma_3 \mathcal{L} = -\mathcal{L} \sigma_3, \quad (2.9)$$

where $\mathcal{L}^*$ has the matrix form $\begin{bmatrix} 0 & -L_+ \\ L_- & 0 \end{bmatrix}$.

We summarize the properties of $\mathcal{L}$ in the following lemma, whose proof is the same as that in [19] and [21, Theorem 2.1]. For convenience of notation, we identify L^2 as a subspace of $\mathbb{C}L^2$ and make no difference between $\psi \in L^2$ and $\mathbf{j}(\psi) \in \mathbb{C}L^2$.

Lemma 2.4 (spectral properties) Suppose the assumptions A0-A1 hold. Let $Q = Q_E$ be a nonlinear ground state, $\|Q_E\|_{L^2} = n$, $0 < n \leq n_0$. Let $\mathcal{L} = \mathcal{L}_E$ be defined as in (2.6).

(1) The eigenvalues of $\mathcal{L}$ are 0 and $\pm i\omega_k$, $k = 1, \dots, N$, where $\omega_k = e_k - e_0 + O(n^2)$ are real and positive. All eigenvalues are simple except 0 which has multiplicity two. The continuous spectrum of $\mathcal{L}$ is

$$\Sigma_c = \{s i : s \in \mathbb{R}, |s| \geq |E|\}. \quad (2.10)$$

There is no embedded eigenvalue. The bottoms of the continuous spectrum, $\pm iE$, are not eigenvalue nor resonance.

(2) The 0-eigenspace is spanned by $\begin{bmatrix} 0 \\ Q \end{bmatrix}$ and $\begin{bmatrix} R \\ 0 \end{bmatrix}$. Note $\mathcal{L} \begin{bmatrix} 0 \\ Q \end{bmatrix} = 0$ and $\begin{bmatrix} R \\ 0 \end{bmatrix}$ is a generalized 0-eigenvector with $\mathcal{L} \begin{bmatrix} R \\ 0 \end{bmatrix} = -\begin{bmatrix} 0 \\ Q \end{bmatrix}$. We denote

$$S = \text{span}_{\mathbb{R}} \left\{ \begin{bmatrix} 0 \\ Q \end{bmatrix}, \begin{bmatrix} R \\ 0 \end{bmatrix} \right\} \subset L^2, \quad (\text{or } S = \text{span}_{\mathbb{R}} \{iQ, R\}).$$

(3) For each eigenvalue $i\omega_k$, $k = 1, \dots, N$, there is an eigenvector Φ_k of the form $\Phi_k = \begin{bmatrix} u_k \\ -iv_k \end{bmatrix}$, where u_k and v_k are real-valued L^2 -functions satisfying

$$L_+ u_k = \omega_k v_k, \quad L_- v_k = \omega_k u_k, \quad (u_k, v_k) = 1.$$

Moreover, they are perturbations of ϕ_k : $u_k, v_k = \phi_k + O(n^2)$. $\bar{\Phi}_k = \begin{bmatrix} u_k \\ iv_k \end{bmatrix}$ is an eigenvector of $-i\omega_k$. We denote the combined eigenspaces of $\pm\omega_k$ as

$$\mathbb{C}\mathbf{E}_k = \text{span}_{\mathbb{C}} \{ \Phi_k, \bar{\Phi}_k \} \subset \mathbb{C}L^2, \quad \mathbf{E}_k = \text{span}_{\mathbb{R}} \left\{ \begin{bmatrix} u_k \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ v_k \end{bmatrix} \right\} \subset L^2.$$

(4) The continuous spectrum subspace, $\mathbf{H}_c(\mathcal{L})$, is equal to

$$\mathbf{H}_c(\mathcal{L}) = \{ \psi \in L^2 : ((\sigma_1 \psi, f)) = 0, \forall f \in S \oplus \mathbf{E}_1 \oplus \dots \oplus \mathbf{E}_N \}.$$

(5) The space $L^2(\mathbb{R}^3, \mathbb{C})$, as a real vector space, can be decomposed as the direct sum of $N + 2$ $\mathcal{L}$ -invariant subspaces:

$$L^2(\mathbb{R}^3, \mathbb{C}) = S \oplus \mathbf{E}_1 \oplus \dots \oplus \mathbf{E}_N \oplus \mathbf{H}_c(\mathcal{L}). \quad (2.11)$$

For any f and g belonging to two different invariant subspaces, we have the orthogonality relation

$$((\sigma_1 f, g)) = 0. \quad (2.12)$$

(6) For any function $\zeta_k \in \mathbf{E}_k$, $k = 1, \dots, N$, there is a unique $\alpha_k \in \mathbb{C}$ so that

$$\zeta_k = \mathbf{R}\mathbf{E} \alpha_k \Phi_k.$$

Since $\mathcal{L} \mathbf{R}\mathbf{E} = \mathbf{R}\mathbf{E} \mathcal{L}$, we have

$$\mathcal{L}\zeta_k = \mathbf{R}\mathbf{E} i\omega_k \alpha_k \Phi_k, \quad e^{t\mathcal{L}}\zeta_k = \mathbf{R}\mathbf{E} e^{ti\omega_k} \alpha_k \Phi_k.$$

(7) By the orthogonality relation (2.12), any $\psi \in L^2$ can be decomposed with respect to (2.11) as

$$\psi = aR + biQ + \sum_{k=1}^N \mathbf{R}\mathbf{E} \alpha_k \Phi_k + \eta, \quad (2.13)$$

with $\eta \in \mathbf{H}_c(\mathcal{L})$, $\alpha_k = c_k + id_k$,

$$\begin{aligned} a &= (Q, R)^{-1}(Q, \operatorname{Re} \psi), & c_k &= (u_k, v_k)^{-1}(v_k, \operatorname{Re} \psi), \\ b &= (Q, R)^{-1}(R, \operatorname{Im} \psi), & d_k &= (u_k, v_k)^{-1}(u_k, \operatorname{Im} \psi). \end{aligned} \quad (2.14)$$

Note $(Q, R)^{-1} = c_1 = O(1)$ and $(u_k, v_k)^{-1} = 1$.

(8) Let $M = \mathbf{E}_1 \oplus \cdots \oplus \mathbf{E}_N \oplus \mathbf{H}_c(\mathcal{L})$. We have $L^2(\mathbb{R}^3, \mathbb{C}) = S \oplus M$ and $M = \begin{bmatrix} Q^\perp \\ R^\perp \end{bmatrix}$. For $m = 0, 1, 2$, there is a constant $C > 1$ such that, for all $\phi \in M \cap H^2$ and all $t \in \mathbb{R}$, we have

$$C^{-1} \|\phi\|_{H^m} \leq \|e^{t\mathcal{L}}\phi\|_{H^m} \leq C \|\phi\|_{H^m}. \quad (2.15)$$

As in [19], in order to prove various estimates and make explicit computations, we will introduce an L^2 -subspace $\mathbf{X}$ and two operators $A : \mathbf{X} \rightarrow \mathbf{X}$ and $U : M \rightarrow \mathbf{X}$ so that

$$\mathcal{L}|_M = U^{-1}(-i)AU. \quad (2.16)$$

Explicitly, let $\mathbf{X}$ be the L^2 -subspace orthogonal to Q :

$$\mathbf{X} = \Pi(L^2) = \{\phi \in L^2(\mathbb{R}^3) : \phi \perp Q\}; \quad \mathbf{X} \longleftrightarrow \begin{bmatrix} Q^\perp \\ Q^\perp \end{bmatrix}, \quad (2.17)$$

where Π is the orthogonal projection which eliminates Q -direction: $\Pi h = h - \frac{(Q, h)}{(Q, Q)} Q$. Let P_M be the projection (not orthogonal) from L^2 onto M according to the decomposition $L^2(\mathbb{R}^3) = S \oplus M$. P_M has the matrix form $\begin{bmatrix} P_1 & 0 \\ 0 & P_2 \end{bmatrix}$, where the projections P_1 and P_2 are given by $(c_1 = (Q, R)^{-1})$

$$\begin{aligned} P_1 : L^2 &\longrightarrow Q^\perp, & P_1 &= \mathbf{id} - c_1 |R\rangle \langle Q|, \\ P_2 : L^2 &\longrightarrow R^\perp, & P_2 &= \mathbf{id} - c_1 |Q\rangle \langle R|. \end{aligned} \quad (2.18)$$

Clearly $P_1 R = 0$ and $P_2 Q = 0$. One can check easily that the maps

$$R_{MX} \equiv \begin{bmatrix} I & 0 \\ 0 & \Pi \end{bmatrix} : M \longrightarrow \mathbf{X}, \quad R_{XM} \equiv \begin{bmatrix} I & 0 \\ 0 & P_2 \end{bmatrix} : \mathbf{X} \longrightarrow M, \quad (2.19)$$

are inverse to each other. We now define $H = L_-$ and

$$A \equiv [(H^2 + H^{1/2} \Pi 2\lambda Q^2 \Pi H^{1/2})]^{1/2} = [H^{1/2} L_+ H^{1/2}]^{1/2}. \quad (2.20)$$

A is a self-adjoint operator acting in $L^2(\mathbb{R}^3)$, with Q as a 0-eigenvector. We shall often view A as an operator restricted to its invariant subspace $\mathbf{X}$. Define

$$U_0 : \mathbf{X} \longrightarrow \mathbf{X}, \quad U_0 \equiv \begin{bmatrix} A^{1/2} H^{-1/2} & 0 \\ 0 & A^{-1/2} H^{1/2} \end{bmatrix}, \quad (2.21)$$

and let

$$U \equiv U_0 R_{MX} : M \longrightarrow \mathbf{X}, \quad U^{-1} \equiv R_{XM} U_0^{-1} : \mathbf{X} \longrightarrow M. \quad (2.22)$$

Notice that $H^{-1/2}$ is defined only in $Q^\perp$. We summarize the properties of A and U in the following lemma.

Lemma 2.5 (Similarity relation) (1) Let $\mathbf{X}$, A , U and U^{-1} be defined as in (2.17), (2.20) and (2.22), respectively. Then (2.16) holds in M . If we denote by P_k^A , $k = 1, \dots, N$, and $\mathbf{P}_c^A$ the orthogonal projections onto the eigenspaces and continuous spectrum subspace of A , we have

$$U P_k^{\mathcal{L}} = P_k^A U P_M, \quad U \mathbf{P}_c^{\mathcal{L}} = \mathbf{P}_c^A U P_M. \quad (2.23)$$

(2) The operators $U : M \rightarrow \mathbf{X}$ and $U^{-1} : \mathbf{X} \rightarrow M$ are bounded in $W^{k,p}$ and L_r^2 norms for $k = 0, 1, 2$, $1 \leq p < \infty$, and $|r| \leq r_1$. Here $r_1 > 0$ is determined by (2.30) later. The operator $U - 1$ is local and bounded by n^2 , and hence so is $[U, i] = [U - 1, i]$, in the sense that

$$\|[U, i]\phi\|_{L^1 \cap L^{5/4}} \leq C n^2 \|\phi\|_{L^5}. \quad (2.24)$$

We have

$$U = U_+ + U_- \text{ conj}, \quad U^{-1} = U_+^* - U_-^* \text{ conj}, \quad (2.25)$$

where **conj** is the conjugation operator with the Pauli matrix σ_3 as its matrix form, and the duals of U_+ and U_- are respect to the L^2 inner product (2.1). The operators U_+ and U_- are given by

$$U_\pm = \frac{1}{2}(\Pi A^{1/2} H^{-1/2} P_1 \pm \Pi A^{-1/2} H^{1/2} \Pi), \quad (2.26)$$

$$(U_\pm)^* = \frac{1}{2}(P_2 H^{-1/2} A^{1/2} \Pi \pm \Pi H^{1/2} A^{-1/2} \Pi). \quad (2.27)$$

In particular, $U_+ = 1 + O(n^2)$, $U_- = O(n^2)$. They are not self-adjoint but they commute with i and **conj**.

(3) For $m = 0, 1, 2$, there is a constant $C > 1$ so that

$$C^{-1} \|\phi\|_{H^m} \leq \|e^{-itA} \phi\|_{H^m} \leq C \|\phi\|_{H^m}, \quad (2.28)$$

for all $\phi \in \mathbf{X} \cap H^m$ and all $t \in \mathbb{R}$. For $q \in [2, \infty]$ and $q' = q/(q-1)$,

$$\|e^{-itA} \mathbf{P}_c^A \Pi \phi\|_{L^q} \leq C |t|^{-3(\frac{1}{2} - \frac{1}{q})} \|\phi\|_{L^{q'}}. \quad (2.29)$$

For sufficiently large $r_1 > 0$ and for all $k, l \in \{1, \dots, N\}$, we have

$$\left\| \langle x \rangle^{-r_1} e^{-itA} \frac{1}{(A - 0i - \omega_k - \omega_l)} \mathbf{P}_c^A \Pi \langle x \rangle^{-r_1} \phi \right\|_{L^2} \leq C \langle t \rangle^{-3/2} \|\phi\|_{L^2}, \quad (2.30)$$

where $0i$ means ri with $\lim_{r \rightarrow 0+}$ outside of the norm. Finally,

$$\begin{aligned} & \left(\phi_0 \phi_k^2, \operatorname{Im} \frac{1}{A - 0i - 2\omega_k} \mathbf{P}_c^A \Pi \phi_0 \phi_k^2 \right) \\ &= \left(\phi_0 \phi_k^2, \operatorname{Im} \frac{1}{H_0 - E - 0i - 2\omega_k} \mathbf{P}_c \phi_0 \phi_k^2 \right) + O(n^2) > 0. \end{aligned} \quad (2.31)$$

Estimate (2.29) for $A = -\Delta + V$ was proven in [9, 24] using estimates from [8, 11]. Estimate (2.30) for $A = (-\Delta + V + m^2)^{1/2}$ was proven in [17]. Lemma 2.5 is a summary of [19, Lemmas 2.5–2.9]. We omit the proof.

3 Stabilization regime

In this section we study the dynamics of the solution when it is close to nonlinear ground states. We want to show that the solution $\psi(t)$ converges to some nonlinear ground state locally as the time tends to infinity. In this time regime, the natural decomposition of $\psi(t)$ is

$$\psi(t) = [Q_E + a(t)R_E + \zeta(t) + \eta(t)] e^{-iEt + i\theta(t)} \quad (3.1)$$

with respect to a fixed E . Here $a(t), \theta(t) \in \mathbb{R}$, $\zeta(t) = \zeta_1(t) + \dots + \zeta_N(t)$, $\zeta_k \in \mathbf{E}_k(\mathcal{L}_E)$, $k = 1, \dots, N$, and $\eta(t) \in \mathbf{H}_c(\mathcal{L}_E)$; see Lemma 2.4. Define

$$\gamma_0^+ \equiv \max_{\substack{1 \leq k, l \leq N \\ |s| < s_0}} \lim_{r \rightarrow 0+} \operatorname{Im} \left(\phi_0 \phi_k \phi_l, \frac{1}{-\Delta + V + e_0 - e_k - e_l - s - ri} \mathbf{P}_c \phi_0 \phi_k \phi_l \right), \quad (3.2)$$

and (recall $c_1 = (Q, R)^{-1}$ and γ_0 is defined in (1.5))

$$D = 6N|c_1|\gamma_0^+/\gamma_0 = O(1). \quad (3.3)$$

We will prove the following theorem.

Theorem 3.1 *Assume the assumptions A0–A3. There are small constant $n_0, \varepsilon_0 > 0$ such that the following holds. Suppose that the initial data ψ_0 with $\|\psi_0\|_{H^1} \ll 1$ is close to a nonlinear ground state $Q_{E_0} e^{i\Theta_0}$ in L_{loc}^2 -norm with $\|Q_{E_0}\|_{L^2} = n \leq n_0$, and that in the decomposition (3.1) of ψ_0 with $E = E_0$ one has*

$$\left(\sum_{k=1}^N \|\zeta_{k, E_0}\|_{L^2}^2 \right)^{1/2} \leq \frac{1}{2}\rho_0, \quad |a_{E_0}| \leq D\rho_0^2, \quad \rho_0 \leq \varepsilon_0 n. \quad (3.4)$$

Suppose, furthermore, for all E close to E_0 with $|E - E_0| \leq 3D\rho_0^2$, the dispersive part $\eta_E(0)$ in the decomposition (3.1) satisfies

$$\begin{aligned} \|e^{s\mathcal{L}} \eta_E(0)\|_{L^5} &\leq n^{4/5} \rho(s)^{8/5}, \\ \|e^{s\mathcal{L}} \eta_E(0)\|_{L_{\text{loc}}^2} &\leq \Lambda(s) \equiv (1+s)^{-1/2} \rho^2(s), \end{aligned} \quad (3.5)$$

for all $s \geq 0$, where

$$\rho(s) = [\rho_0^{-2} + N^{-1}\gamma_0 n^2 s]^{-1/2}. \quad (3.6)$$

Then there is a frequency E_∞ with $|E_\infty - E_0| \leq 3D\rho_0^2$ and a function $\Theta(t) = -E_\infty t + O(\log(t))$ for $t \in [0, \infty)$ such that, for some constant $C_3 > 1$ independent of n ,

$$\|\psi(t) - Q_{E_\infty} e^{i\Theta(t)}\|_{L_{\text{loc}}^2} \leq C_3 \rho(t), \quad (t \geq 0). \quad (3.7)$$

Suppose, furthermore, that $\left(\sum_{k=1}^N \|\zeta_{k,E_0}\|_{L^2}^2\right)^{1/2} \geq \frac{1}{4}\rho_0$. Then we also have a lower bound

$$C_3^{-1} \rho(t) \leq \|\psi(t) - Q_{E_\infty} e^{i\Theta(t)}\|_{L_{\text{loc}}^2}, \quad (t \geq 0). \quad (3.8)$$

We call Eq. (3.5) the *out-going estimates* of $\eta(0)$. The theorem holds true if they are replaced by the following stronger but simpler assumption that

$$\psi_0 \in Y \equiv H^1 \cap L^1(\mathbb{R}^3), \quad \|\eta(0)\|_Y \leq \rho_0^2, \quad (3.9)$$

since (3.9) implies (3.5) by Lemma 2.3. Eq. (3.9) means that the data ψ_0 is localized. In contrast, Eq. (3.5) only requires the data to be “out-going” in some sense. This will be useful when we prove Theorem 1.1 using Theorem 3.1 in section 4.

Our strategy of proof is as follows. For each $T > 0$, we choose $Q_{E(T)}$ to be the best approximation of $\psi(T)$ given by Lemma 2.2. We will prove estimates for the components of $\psi(t)$ in the decomposition (3.1) with respect to $E = E(T)$ for $t \in [0, T]$. We will then use a continuity argument to show that $E(T)$ can always be chosen and we have uniform estimates as $T \rightarrow \infty$. It then follows that $E(T)$ converges to some E_∞ close to $E(0)$ as $T \rightarrow \infty$.

3.1 Equations

Let $Q = Q_E$ be a fixed nonlinear ground state with frequency E near e_0 , and $R = R_E = \partial_E Q_E$. We write the solution $\psi(t, x)$ of (1.1) in the form

$$\psi(t, x) = [Q_E(x) + a(t)R_E(x) + h(t, x)] e^{-iEt + i\theta(t)}, \quad (3.10)$$

where $a(t), \theta(t) \in \mathbb{R}$ and $h(t, \cdot) \in M_E$. M_E is defined in Lemma 2.4 (8). Substituting the ansatz (3.10) into (1.1) and using $\mathcal{L}iQ = 0$ and $\mathcal{L}R = -iQ$, we get

$$\partial_t h = \mathcal{L}h + i^{-1}(F + \dot{\theta}(Q + aR + h)) - aiQ - \dot{a}R. \quad (3.11)$$

Here

$$F = \lambda Q(2|h_a|^2 + h_a^2) + \lambda|h_a|^2 h_a, \quad h_a = aR + h.$$

We want to choose $a(t)$ and $\theta(t)$ so that $h(t) \in M$, that is, $h(0) \in M$ and $i^{-1}(F + \dot{\theta}(Q + aR + h)) - aiQ - \dot{a}R \in M$. Since $M = \begin{bmatrix} Q^\perp \\ R^\perp \end{bmatrix}$, $a(t)$ and $\theta(t)$ satisfy

$$\begin{aligned} \left(Q, \text{Im}(F + \dot{\theta}h) - \dot{a}R \right) &= 0, \\ \left(R, \text{Re}(F + \dot{\theta}(Q + aR + h)) - aQ \right) &= 0. \end{aligned}$$

Denote $c_1 = (Q, R)^{-1}$. We have

$$\begin{aligned} \dot{a} &= (c_1 Q, \text{Im}(F + \dot{\theta}h)), \\ \dot{\theta} &= -[a + (c_1 R, \text{Re } F)] \cdot [1 + a(c_1 R, R) + (c_1 R, \text{Re } h)]^{-1}. \end{aligned}$$

Eq. (3.11) for h becomes

$$\partial_t h = \mathcal{L}h + P_M F_{all}, \quad F_{all} = i^{-1}(F + \dot{\theta}(aR + h)). \quad (3.12)$$

We decompose $h(t)$ with respect to the spectral decomposition (2.11),

$$h(t) = \zeta(t) + \eta(t), \quad \zeta = \zeta_1 + \cdots + \zeta_N,$$

where $\zeta_k \in \mathbf{E}_k(\mathcal{L})$, $k = 1, \dots, N$, and $\eta \in \mathbf{H}_c(\mathcal{L})$. For each $\zeta_k(t)$ we associate a function $z_k(t) \in \mathbb{C}$ by writing $\zeta_k = \text{Re}(z_k) u_k + \text{Im}(z_k) i v_k$. In other words, $\zeta_k(t) = \mathbf{R} \mathbf{E} z_k(t) \Phi_k$. If we define $u_k^\pm = \frac{1}{2}(u_k \pm v_k)$, we can write

$$\zeta_k = \text{Re}(z_k) u_k + \text{Im}(z_k) i v_k = z_k u_k^+ + \bar{z}_k u_k^-. \quad (3.13)$$

Projecting (3.12) to $\mathbf{E}_k(\mathcal{L})$ we get

$$\dot{z}_k = -i\omega_k z_k + (v_k, \text{Re } F_{all}) + i(u_k, \text{Im } F_{all}) = -i\omega_k z_k + (u_k^+, F_{all}) - (u_k^-, \bar{F}_{all}).$$

From the linear part, we identify the *phase factor* of z_k as $-\omega_k$. Hence we define $p_k(t) = e^{i\omega_k t} z_k(t)$, which has the same magnitude as z_k but with no strong oscillation. $p_k(t)$ satisfy

$$\begin{aligned} e^{-i\omega_k t} \dot{p}_k(t) &= (u_k^+, F_{all}) - (u_k^-, \bar{F}_{all}) \\ &= i^{-1} \left\{ (u_k^+, F) + (u_k^-, \bar{F}) + [(u_k^+, h) + (u_k^-, \bar{h}) + (u_k, R)a] \dot{\theta} \right\}. \end{aligned}$$

Also, projecting (3.12) to $\mathbf{H}_c(\mathcal{L})$ we get $\partial_t \eta = \mathcal{L} \eta + \mathbf{P}_c^\mathcal{L} F_{all}$.

Summarizing, for

$$\begin{aligned} \psi(t) &= (Q + a(t)R + h(t)) e^{-iEt + i\theta(t)}, \\ h &= \zeta + \eta, \quad \zeta = \zeta_1 + \cdots + \zeta_N, \quad \zeta_k = z_k u_k^+ + \bar{z}_k u_k^-, \quad z_k = e^{-i\omega_k t} p_k, \end{aligned}$$

we have

$$\begin{aligned}\dot{a} &= (c_1 Q, \operatorname{Im}(F + \dot{\theta} h)), \\ i e^{-i\omega_k t} \dot{p}_k &= (u_k^+, F) + (u_k^-, \bar{F}) + [(u_k^+, h) + (u_k^-, \bar{h}) + (u_k, R)a] \dot{\theta}, \\ \partial_t \eta &= \mathcal{L} \eta + \mathbf{P}_c^\mathcal{L} i^{-1} (F + \dot{\theta} (aR + h)),\end{aligned}\tag{3.14}$$

where $c_1 = (Q, R)^{-1}$,

$$F = \lambda Q(2|h_a|^2 + h_a^2) + \lambda |h_a|^2 h_a, \quad h_a = aR + h, \tag{3.15}$$

$$\dot{\theta} = F_\theta \equiv -[a + (c_1 R, \operatorname{Re} F)] \cdot [1 + a(c_1 R, R) + (c_1 R, \operatorname{Re} h)]^{-1}. \tag{3.16}$$

This is a system of equations involving a , z_k and η only. Note that θ enters (3.14) only via $\dot{\theta} = F_\theta$. It will appear in the form $e^{i\theta}$ when we integrate η . Hence we do not need estimates of θ for the proof.

For convenience, we will use the following convention.

Convention 3.1 For $k = 1, 2, \dots, N$, denote

$$\omega_{-k} = -\omega_k, \quad z_{-k} = \bar{z}_k, \quad p_{-k} = \bar{p}_k. \tag{3.17}$$

We have $z_k(t) = e^{-i\omega_k t} p_k(t)$ for both $k > 0$ and $k < 0$. We also denote

$$\Omega = \{\pm 1, \dots, \pm N\}. \tag{3.18}$$

3.2 Decompositions of F , a and η

Most quantities in our system of equations are strongly oscillatory. It is necessary to identify their oscillatory parts before we can estimate. In this subsection we identify the leading oscillatory terms of a and η , and decompose F according to order. We will treat z_k and a again in §3.3. Note that

$$Q = O(n), \quad R = O(n^{-1}), \quad c_1 = O(1), \quad u_k^+ = \phi_k + O(n^2), \quad u_k^- = O(n^2). \tag{3.19}$$

In fact, since $\|Q_{E_0}\|_{L^2} = n$ and $|E - E_0| \leq 3D\varepsilon_0^2 n^2$, we have $\|Q_E\|_{L^2} = [1 + O(\varepsilon_0^2)]n$ and $Q = (1 + O(\varepsilon_0^2))n\phi_0 + O(n^3)$. We will prove that

$$|z_k(t)| \leq Ct^{-1/2}, \quad |a(t)| + \|\eta(t)\|_{L_{\text{loc}}^2} \leq Ct^{-1}, \quad \text{as } t \rightarrow \infty. \tag{3.20}$$

Hence the main term in $h_a = aR + \zeta + \eta$ is ζ . Therefore, the main part of F , defined in (3.15), is

$$F_1 = \lambda Q(2|\zeta|^2 + \zeta^2). \tag{3.21}$$

3.2.1 Decomposition of a

We now identify the main oscillatory terms of $a(t)$. Recall from (3.14) that $\dot{a} = (c_1 Q, \text{Im } F + \dot{\theta} h)$, $c_1 = (Q, R)^{-1}$. We shall impose the boundary condition of a at $t = T$, which is in fact a condition imposed on the choice of $E(T)$. Hence we use the following equivalent integral equation:

$$a(t) = a(T) + \int_T^t (c_1 Q, \text{Im } F + \dot{\theta} h)(s) ds.$$

The main term of $\text{Im}(F + \dot{\theta} h)$ is $\text{Im } F_1 = \text{Im } \lambda Q \zeta^2$. Thus the main oscillatory terms of $a(t)$ are from the integral $\int_T^t A^{(2)} ds$ with

$$A^{(2)} \equiv (c_1 Q, \text{Im } \lambda Q \zeta^2) = (c_1 \lambda Q^2, \text{Im } \sum_{k,l=1}^N \zeta_k \zeta_l).$$

Since

$$\begin{aligned} \text{Im } \zeta_k \zeta_l &= \text{Im}(z_k u_k^+ + \bar{z}_k u_k^-)(z_l u_l^+ + \bar{z}_l u_l^-) \\ &= \text{Im}(z_k z_l)(u_k^+ u_l^+ - u_k^- u_l^-) + \text{Im}(z_k \bar{z}_l)(u_k^+ u_l^- - u_k^- u_l^+), \end{aligned}$$

we have

$$A^{(2)} = \sum_{k,l=1}^N \{a_{kl,1} \text{Im}(z_k z_l) + a_{kl,2} \text{Im}(z_k \bar{z}_l)\},$$

where $a_{kl,1} = (c_1 \lambda Q^2, (u_k^+ u_l^+ - u_k^- u_l^-))$ and $a_{kl,2} = (c_1 \lambda Q^2, u_k^+ u_l^- - u_k^- u_l^+)$ are real constants bounded by n^2 . We can integrate by parts $A^{(2)}$ to get:

$$\begin{aligned} \int_T^t A^{(2)} ds &= \text{Im} \sum_{k,l=1}^N \int_T^t a_{kl,1}(z_k z_l) + a_{kl,2}(z_k \bar{z}_l) ds \\ &= \text{Im} \sum_{k,l=1}^N \int_T^t e^{-i(\omega_k + \omega_l)s} a_{kl,1}(p_k p_l) + e^{-i(\omega_k - \omega_l)s} a_{kl,2}(p_k \bar{p}_l) ds \\ &= \text{Im} \sum_{k,l=1}^N [i a_{kl,3} z_k z_l + i a_{kl,4} z_k \bar{z}_l]_T^t - \int_T^t A_{2,rm d}(s) ds, \end{aligned} \quad (3.22)$$

where

$$a_{kl,3} = (\omega_k + \omega_l)^{-1} a_{kl,1}, \quad a_{kl,4} = \delta_k^l (\omega_k - \omega_l)^{-1} a_{kl,2}, \quad (3.23)$$

$$A_{2,rm d} = \text{Im} \sum_{k,l=1}^N \left\{ e^{-i(\omega_k + \omega_l)s} i a_{kl,3} \frac{d}{ds} (p_k p_l) + e^{-i(\omega_k - \omega_l)s} i a_{kl,4} \frac{d}{ds} (p_k \bar{p}_l) \right\}. \quad (3.24)$$

Here we put δ_k^l in the definition of $a_{kl,4}$ to impose $a_{kk,4} = 0$. We have the factor δ_k^l in $a_{kl,4}$ since $\text{Im } z_k \bar{z}_l = 0$ if $k = l$. Also note that $a_{kl,3}$ and $a_{kl,4}$ are real constants bounded by n^2 . The first part in (3.22) can be rewritten as $a^{(2)}(t) - a^{(2)}(T)$, where

$$a^{(2)} = \text{Im} \sum_{k,l=1}^N i a_{kl,3} z_k z_l + i a_{kl,4} z_k \bar{z}_l = \sum_{k,l \in \Omega} a_{kl} z_k z_l. \quad (3.25)$$

Here we have used Convention 3.1. The constants $a_{kl} = \frac{1}{2} a_{kl,3}$ if k and l have the same sign; $a_{kl} = \frac{1}{2} a_{kl,4}$ otherwise. In particular, a_{kl} are real constants bounded by n^2 . $a^{(2)}(t)$ contains the main oscillatory part of a . We denote the rest of $a(t)$ by $b(t)$,

$$a(t) = a^{(2)}(t) + b(t). \quad (3.26)$$

Thus $b(t) = a(T) - a^{(2)}(T) + \int_T^t \dot{b}(s) ds$ with

$$\dot{b} = (c_1 Q, \text{Im}[F - F_1 + \dot{\theta} h]) - A_{2, \text{rmd}}. \quad (3.27)$$

As we will see later that $\dot{b}(t)$ is smaller than $\dot{a}^{(2)}(t)$. However, $b(t)$ is the main part of $a(t)$: We have $b(t) \lesssim \rho^2(t)$ while $a^{(2)}(t) \lesssim n^2 \rho^2(t)$.

3.2.2 Decompositions of F

Recall (3.15),

$$F = \lambda Q(2|h_a|^2 + h_a^2) + \lambda|h_a|^2 h_a, \quad h_a = aR + h.$$

In view of (3.20), we decompose $h_a = \zeta + bR + (\eta + a^{(2)}R)$ and decompose F as

$$F = F_1 + F_2 + F_3 + F_4 + F_5, \quad (3.28)$$

where

$$\begin{aligned} F_1 &= \lambda Q(2|\zeta|^2 + \zeta^2), \\ F_2 &= 2\lambda Q R b(2\zeta + \bar{\zeta}) + 3\lambda Q R^2 b^2 + \lambda(\zeta + bR)^2(\bar{\zeta} + bR), \\ F_3 &= 2\lambda Q R a^{(2)}(2\zeta + \bar{\zeta}), \\ F_4 &= 2\lambda Q[(\zeta + \bar{\zeta})\eta^{(2)} + \zeta \bar{\eta}^{(2)}], \\ F_5 &= 2\lambda Q[(\zeta + \bar{\zeta})\eta^{(3)} + \zeta \bar{\eta}^{(3)}] \\ &\quad + \lambda Q[2|\eta_a|^2 + \eta_a^2] + 2\lambda Q R b(2\eta_a + \bar{\eta}_a) \quad (\eta_a = \eta + a^{(2)}R) \\ &\quad + \lambda(aR + h)^2(aR + \bar{h}) - \lambda(\zeta + bR)^2(\bar{\zeta} + bR). \end{aligned} \quad (3.29)$$

Here F_1 consists of terms of order nz^2 ; F_2 , F_3 and F_4 consist of terms no smaller than $n^2 z^3$; and F_5 higher order terms.

3.2.3 Decomposition of η

We now identify the main terms in η . We first recall from (3.14) that

$$\partial_t \eta = \mathcal{L} \eta + \mathbf{P}_c^\mathcal{L} i^{-1} [F + \dot{\theta}(aR + \zeta + \eta)].$$

Using Lemma 2.5 that $\mathcal{L} = U^{-1}(-i)AU$ on $\mathbf{H}_c(\mathcal{L})$ and $U \mathbf{P}_c^\mathcal{L} = \mathbf{P}_c^A U$, we have

$$\begin{aligned} \partial_t U \eta &= -iAU \eta + \mathbf{P}_c^A U i^{-1} [F + \dot{\theta}(aR + \zeta + \eta)] \\ &= -iAU \eta - i\dot{\theta}U \eta + \mathbf{P}_c^A U i^{-1} [F + \dot{\theta}(aR + \zeta)] - \mathbf{P}_c^A [U, i] \dot{\theta} \eta. \end{aligned}$$

Here we have used the commutator $[U, i]$ to interchange U and i so as to produce the term $i\dot{\theta}U \eta$. This term is a global linear term in $U \eta$ and cannot be treated as error (however $[U, i]U^{-1}\dot{\theta}\eta$ is a local error term). We can eliminate it by introducing

$$\tilde{\eta}(t) \equiv e^{i\theta(t)} U \eta(t), \quad \theta(t) = \int_0^t F_\theta(s) ds. \quad (3.30)$$

We have $\tilde{\eta}(0) = U \eta(0)$ and

$$\partial_t \tilde{\eta} = -iA \tilde{\eta} + e^{i\theta} \mathbf{P}_c^A U i^{-1} [F + \dot{\theta}(aR + \zeta)] - e^{i\theta} \mathbf{P}_c^A [U, i] \dot{\theta} \eta.$$

Hence $\tilde{\eta}(t)$ satisfies the integral equation (using $e^{-iAt} \tilde{\eta}(0) = U e^{t\mathcal{L}} \eta(0)$)

$$\tilde{\eta}(t) = U e^{t\mathcal{L}} \eta(0) + \int_0^t e^{-iA(t-s)} \mathbf{P}_c^A F_\eta(s) ds, \quad (3.31)$$

$$F_\eta \equiv e^{i\theta} U i^{-1} [F + \dot{\theta}(aR + \zeta)] - e^{i\theta} [U, i] \dot{\theta} \eta. \quad (3.32)$$

Since U and U^{-1} are bounded in Sobolev spaces by Lemma 2.5, and

$$\eta(t) = U^{-1} e^{-i\theta(t)} \tilde{\eta}(t), \quad (3.33)$$

for the purpose of estimation we can treat η and $\tilde{\eta}$ as the same.

To identify the main term of $\tilde{\eta}$, we decompose F_η as follows,

$$\begin{aligned} F_\eta &= F_{\eta,2} + F_{\eta,3}, \\ F_{\eta,2} &= e^{i\theta} U i^{-1} F_1, \\ F_{\eta,3} &= e^{i\theta} U i^{-1} [(F - F_1) + \dot{\theta}(aR + \zeta)] - e^{i\theta} [U, i] \dot{\theta} \eta. \end{aligned} \quad (3.34)$$

The leading part of $\tilde{\eta}$ is from $F_{\eta,2}$. Recall $F_1 = \lambda Q(\zeta^2 + 2\zeta\bar{\zeta})$, and $U = U_+ + \mathbf{conj} U_-$ with U_+ and U_- commuting with i and $\mathbf{conj}$, see (2.25). Hence

$$\begin{aligned} F_{\eta,2} &= e^{i\theta} U i^{-1} \lambda Q(\zeta^2 + 2\zeta\bar{\zeta}) \\ &= e^{i\theta} i^{-1} (U_+ - \mathbf{conj} U_-) \lambda Q(\zeta^2 + 2\zeta\bar{\zeta}) \\ &= e^{i\theta} i^{-1} \{ U_+ \lambda Q(\zeta^2 + 2\zeta\bar{\zeta}) - U_- \lambda Q(\bar{\zeta}^2 + 2\bar{\zeta}\zeta) \}. \end{aligned}$$

Substituting $\zeta = \sum_{l=1}^N \zeta_l$ and $\zeta_l = z_l u_l^+ + \bar{z}_l u_l^-$, we have

$$\begin{aligned} F_{\eta,2} &= e^{i\theta} i^{-1} \sum_{k,l=1}^N \{U_+ \lambda Q(\zeta_k \zeta_l + \zeta_k \bar{\zeta}_l + \bar{\zeta}_k \zeta_l) - U_- \lambda Q(\bar{\zeta}_k \bar{\zeta}_l + \zeta_k \bar{\zeta}_l + \bar{\zeta}_k \zeta_l)\} \\ &= e^{i\theta} i^{-1} \sum_{k,l \in \Omega} z_k z_l \Phi_{kl}. \end{aligned}$$

In the last line, Convention 3.1 is used. In particular, $z_k = \bar{z}_{|k|}$ if $k < 0$. The functions Φ_{kl} are defined as follows. For $k < 0$, denote $u_k^+ = u_{|k|}^+$ and $u_k^- = u_{|k|}^-$. We define

$$\begin{aligned} k, l > 0 : \quad \Phi_{kl} &= U_+ \lambda Q(u_k^+ u_l^+ + u_k^+ u_l^- + u_k^- u_l^+) \\ &\quad - U_- \lambda Q(u_k^- u_l^- + u_k^+ u_l^- + u_k^- u_l^+), \\ k > 0, l < 0 : \quad \Phi_{kl} &= U_+ \lambda Q(u_k^+ u_l^- + u_k^+ u_l^+ + u_k^- u_l^-) \\ &\quad - U_- \lambda Q(u_k^- u_l^+ + u_k^+ u_l^+ + u_k^- u_l^-), \\ k < 0, l > 0 : \quad \Phi_{kl} &= \Phi_{lk}, \\ k, l < 0 : \quad \Phi_{kl} &= U_+ \lambda Q(u_k^- u_l^- + u_k^+ u_l^- + u_k^- u_l^+) \\ &\quad - U_- \lambda Q(u_k^+ u_l^+ + u_k^+ u_l^- + u_k^- u_l^+). \end{aligned} \tag{3.35}$$

Since $u_k^- = O(n^2)$ and $U_- = O(n^2)$, we have $\Phi_{kl} = O(n^3)$ if $k, l < 0$, $\Phi_{kl} = O(n)$ otherwise.

We now integrate $F_{\eta,2}$. Using $z_k(s) = e^{-i\omega_k s} p_k(s)$ (see Convention 3.1) and integrating by parts we get

$$\begin{aligned} \int_0^t e^{-iA(t-s)} \mathbf{P}_c^A F_{\eta,2} ds &= \sum \int_0^t e^{-iA(t-s)} e^{i\theta} i^{-1} z_k z_l \mathbf{P}_c^A \Phi_{kl} ds \\ &= \sum i^{-1} \int_0^t e^{-iAt} e^{is(A-0i-\omega_k-\omega_l)} (e^{i\theta} p_k p_l) \mathbf{P}_c^A \Phi_{kl} ds \\ &= \sum e^{i\theta} z_k z_l \tilde{\eta}_{kl} - \sum e^{-iAt} (e^{i\theta} z_k z_l)(0) \tilde{\eta}_{kl} \\ &\quad - \sum \int_0^t e^{-iAt} e^{is(A-0i-\omega_k-\omega_l)} \frac{d}{ds} (e^{i\theta} p_k p_l(s)) \tilde{\eta}_{kl} ds, \end{aligned}$$

where the summation is over $k, l \in \Omega$ and

$$\tilde{\eta}_{kl} = \lim_{r \rightarrow 0+} \frac{-1}{A - \omega_k - \omega_l - ri} \mathbf{P}_c^A \Phi_{kl}, \quad (k, l \in \Omega). \tag{3.36}$$

Here we add $-ri$ since $A - \omega_k - \omega_l$ may not be invertible. We take $r \rightarrow 0+$, not $r \rightarrow 0-$, to get the decay of the second and the last terms above, see Lemma 2.5 (3). Recall $\omega_k \sim e_k - e_0 > 0$ for $k > 0$ and $\omega_k = -\omega_{|k|}$ if $k < 0$. Since A has a spectral gap $\sim |e_0|$, $A - \omega_k - \omega_l$ is not invertible only if $k > 0$ and $l > 0$, by Assumption A2. Hence $\tilde{\eta}_{kl} \in L^2$ unless both k and l are positive.

We denote the main term above by $\tilde{\eta}^{(2)}$,

$$\tilde{\eta}^{(2)} = \sum_{k,l \in \Omega} e^{i\theta} z_k z_l \tilde{\eta}_{kl}, \tag{3.37}$$

and denote the rest of $\tilde{\eta}$ by $\tilde{\eta}^{(3)}$. We decompose $\tilde{\eta}^{(3)} = \tilde{\eta}_1^{(3)} + \cdots + \tilde{\eta}_4^{(3)}$ where

$$\begin{aligned}\tilde{\eta}_1^{(3)} &= U e^{t\mathcal{L}} \eta(0), \\ \tilde{\eta}_2^{(3)} &= - \sum_{k,l \in \Omega} e^{-iAt} (e^{i\theta} z_k z_l)(0) \tilde{\eta}_{kl} = -e^{-iAt} \tilde{\eta}^{(2)}(0), \\ \tilde{\eta}_3^{(3)} &= - \sum_{k,l \in \Omega} \int_0^t e^{-iAt} e^{is(A-0i-\omega_k-\omega_l)} \frac{d}{ds} (e^{i\theta} p_k p_l(s)) \tilde{\eta}_{kl} ds, \\ \tilde{\eta}_4^{(3)} &= \int_0^t e^{-iA(t-s)} \mathbf{P}_c^A F_{\eta,3} ds.\end{aligned}\tag{3.38}$$

We have

$$\tilde{\eta} = \tilde{\eta}^{(2)} + \tilde{\eta}^{(3)}, \quad \tilde{\eta}^{(3)} = \tilde{\eta}_1^{(3)} + \cdots + \tilde{\eta}_4^{(3)}.$$

Correspondingly, we define

$$\begin{aligned}\eta^{(2)} &= U^{-1} e^{-i\theta} \tilde{\eta}^{(2)} = U^{-1} \sum_{k,l \in \Omega} z_k z_l \tilde{\eta}_{kl}, \\ \eta^{(3)} &= U^{-1} e^{-i\theta} \tilde{\eta}^{(3)}, \quad \eta_j^{(3)} = U^{-1} e^{-i\theta} \tilde{\eta}_j^{(3)}, \quad (j = 1, \dots, 4).\end{aligned}\tag{3.39}$$

Lemma 3.2 *Suppose, for a fixed time, for some $\rho \ll n \leq n_0$,*

$$\begin{aligned}\|Q\| &= n, \quad \|\eta\|_{L^2 \cap L^5} \ll 1, \quad \|\eta\|_{L_{\text{loc}}^2} \leq n, \\ \max_{k=1, \dots, N} |z_k| &\leq \rho, \quad |a| \leq C\rho^2.\end{aligned}\tag{3.40}$$

Denote

$$\begin{aligned}\tilde{X} &= \rho^2 \|\eta\|_{L_{\text{loc}}^2}^2 + n \|\eta\|_{L_{\text{loc}}^2}^2 + \|\eta^3\|_{L_{\text{loc}}^1}, \\ X &= n\rho \|\eta\|_{L_{\text{loc}}^2}^2 + n \|\eta\|_{L_{\text{loc}}^2}^2 + \|\eta^3\|_{L_{\text{loc}}^1}.\end{aligned}\tag{3.41}$$

We have

$$\begin{aligned}\|F_5\|_{L_{\text{loc}}^1} &\lesssim n\rho^4 + n\rho \|\eta^{(3)}\|_{L_{\text{loc}}^2} + \tilde{X}, \\ \|F_3 + F_4 + F_5\|_{L_{\text{loc}}^1} &\lesssim n^2 \rho^3 + X, \\ \|F - F_1\|_{L_{\text{loc}}^1} &\lesssim \rho^3 + X, \quad \|F\|_{L_{\text{loc}}^1} \lesssim n\rho^2 + X, \\ \|F_\theta\|_{L_{\text{loc}}^1} &\lesssim \rho^2 + n^{-1}X, \\ \max_k |\dot{p}_k| &\lesssim n\rho^2 + X, \quad |\dot{b}| \lesssim n\rho^3 + nX, \\ \|F_{\eta,3}\|_{L^{5/4} \cap L^1} &\lesssim \rho^3 + n\rho \|\eta\|_{L_{\text{loc}}^2} + n \|\eta\|_{L_{\text{loc}}^2 \cap L^5}^2 + \|\eta^3\|_{L^{5/4} \cap L^1} + n^2 \rho^2 \|\eta\|_{L^5}, \\ \|F_\eta\|_{L^{5/4}} &\lesssim n\rho^2 + n\rho \|\eta\|_{L_{\text{loc}}^2} + n \|\eta\|_{L_{\text{loc}}^2 \cap L^5}^2 + \|\eta^3\|_{L^{5/4}} + n^2 \rho^2 \|\eta\|_{L^5}.\end{aligned}\tag{3.42}$$

PROOF: By (3.40) we have $|a^{(2)}| \lesssim n^2 \rho^2$, $|b| \lesssim \rho^2$ and $\|\eta^{(2)}\|_{L^2_{\text{loc}}} \leq n\rho^2$. Also recall (3.19). Hence for F_5 defined in (3.29) we have

$$\begin{aligned} \|F_5\|_{L^1_{\text{loc}}} &\lesssim n\rho \|\eta^{(3)}\|_{L^2_{\text{loc}}} + n \left(\|\eta\|_{L^2_{\text{loc}}} + n\rho^2 \right)^2 + \rho^2 \left(\|\eta\|_{L^2_{\text{loc}}} + n\rho^2 \right) + \|\eta^3\|_{L^1_{\text{loc}}} \\ &\lesssim n\rho^4 + n\rho \|\eta^{(3)}\|_{L^2_{\text{loc}}} + \rho^2 \|\eta\|_{L^2_{\text{loc}}} + n \|\eta\|_{L^2_{\text{loc}}}^2 + \|\eta^3\|_{L^1_{\text{loc}}} \\ &= n\rho^4 + n\rho \|\eta^{(3)}\|_{L^2_{\text{loc}}} + \tilde{X}, \end{aligned}$$

where $\tilde{X}$ is defined in (3.41). Combining $\eta^{(2)} + \eta^{(3)} = \eta$ in the definition (3.29) of F_4 and F_5 , we can use the same argument for $\|F_5\|_{L^1_{\text{loc}}}$ to show

$$\|F_4 + F_5\|_{L^1_{\text{loc}}} \lesssim n\rho^4 + n\rho \|\eta\|_{L^2_{\text{loc}}} + \tilde{X} \lesssim n\rho^4 + X.$$

By their explicit form in (3.29), we have $\|F_3\|_{L^1_{\text{loc}}} \lesssim n^2 \rho^3$, $\|F_2\|_{L^1_{\text{loc}}} \sim |bz| + b^2/n + (z + b/n)^3 \lesssim \rho^3$, and $\|F_1\|_{L^1_{\text{loc}}} \lesssim n\rho^2$. Hence

$$\|F_3 + F_4 + F_5\|_{L^1_{\text{loc}}} \lesssim n^2 \rho^3 + X, \quad \|F - F_1\|_{L^1_{\text{loc}}} \lesssim \rho^3 + X, \quad \|F\|_{L^1_{\text{loc}}} \lesssim n\rho^2 + X.$$

From definition (3.16) of F_θ we have

$$|F_\theta| \lesssim |a| + n^{-1} \|F\|_{L^1_{\text{loc}}} \lesssim \rho^2 + n^{-1}(n\rho^2 + X) \lesssim \rho^2 + n^{-1}X.$$

By (3.14), we have

$$\begin{aligned} |\dot{p}_k| &\lesssim \|F\|_{L^1_{\text{loc}}} + (\rho + \|\eta\|_{L^2_{\text{loc}}})|F_\theta| \\ &\lesssim n\rho^2 + X + (\rho + \|\eta\|_{L^2_{\text{loc}}})(\rho^2 + n^{-1}X) \lesssim n\rho^2 + X. \end{aligned}$$

By (3.27),

$$|\dot{b}| \leq Cn \|F - F_1\|_{L^1_{\text{loc}}} + |(c_1 Q, \text{Im } h)| \cdot |F_\theta| + |A_{2,\text{rmd}}|.$$

Note that $|(c_1 Q, \text{Im } h)| \lesssim n^3 \|h\|_{L^1_{\text{loc}}}$ since $\text{Im } h \perp R$ and $Q = Cn^2 R + O(n^3)$ by Lemma 2.1. Also $|A_{2,\text{rmd}}| \lesssim n^2 \rho \max_k |\dot{p}_k|$ by definition (3.24) of $A_{2,\text{rmd}}$. Thus

$$|\dot{b}| \lesssim n(\rho^3 + X) + n^3(\rho + \|\eta\|_{L^2_{\text{loc}}})(\rho^2 + n^{-1}X) + n^2 \rho(n\rho^2 + X) \lesssim n\rho^3 + nX.$$

Let $r = 1$ or $r = 5/4$. The estimate for $\|F - F_1\|_{L^r}$ is the same as that for $\|F - F_1\|_{L^1_{\text{loc}}}$ except for the non-local term $\lambda \eta^2 \bar{\eta}$ and $\|n\eta^2\|_{L^r} \lesssim n \|\eta\|_{L^2_{\text{loc}} \cap L^5}^2$. Thus

$$\|F - F_1\|_{L^r} \lesssim \rho^3 + n\rho \|\eta\|_{L^2_{\text{loc}}} + n \|\eta\|_{L^2_{\text{loc}} \cap L^5}^2 + \|\eta^3\|_{L^r}.$$

From the definition (3.34) of $F_{\eta,3}$, the boundedness of U , and the estimate (2.24) for $[U, i]$, we have

$$\begin{aligned} \|F_{\eta,3}\|_{L^r} &\lesssim \|F - F_1\|_{L^r} + |F_\theta|(n^{-1}|a| + \max |z_k|) + n^2 |F_\theta| \|\eta\|_{L^5} \\ &\lesssim \rho^3 + n\rho \|\eta\|_{L^2_{\text{loc}}} + n \|\eta\|_{L^2_{\text{loc}} \cap L^5}^2 + \|\eta^3\|_{L^r} \\ &\quad + (\rho^2 + n^{-1}X)(\rho + n^2 \|\eta\|_{L^5}) \\ &\lesssim \rho^3 + n\rho \|\eta\|_{L^2_{\text{loc}}} + n \|\eta\|_{L^2_{\text{loc}} \cap L^5}^2 + \|\eta^3\|_{L^r} + n^2 \rho^2 \|\eta\|_{L^5}. \end{aligned}$$

Here we also used $\|\eta^3\|_{L^1_{\text{loc}}} \lesssim \|\eta^3\|_{L^r}$. Since $\|F_\eta\|_{L^{5/4}} \lesssim n\rho^2 + \|F_{\eta,3}\|_{L^{5/4}}$, the estimate for $\|F_\eta\|_{L^{5/4}}$ in (3.42) follows. **Q.E.D.**

3.3 Normal forms for equations of bound states

Recall that $z_k(t) = e^{-i\omega_k t} p_k(t)$ and $a(t) = a^{(2)}(t) + b(t)$. Since many terms in the equations of p_k and b are oscillatory and only contribute to the phases, we now derive the normal forms for the equations of p_k and b , where those terms are removed. Throughout this subsection, Convention 3.1 for ω_k , z_k and p_k with $k < 0$ is understood.

Lemma 3.3 (Normal Form) *Suppose, for some $\rho(t) \ll n \leq n_0$,*

$$\begin{aligned} \|Q\| = n, \quad \|\eta(t)\|_{L^2 \cap L^5} \ll 1, \quad \|\eta(t)\|_{L^2_{\text{loc}}} \leq n, \\ \max_{k=1, \dots, N} |z_k(t)| \leq \rho(t), \quad |a(t)| \leq C\rho^2(t). \end{aligned} \quad (3.43)$$

There are perturbations $q_k(t)$ of $p_k(t)$, $k = 1, \dots, N$, and $\beta(t)$ of $b(t)$, to be defined in (3.96) and (3.103), satisfying

$$|q_k - p_k| \leq Cn\rho^2, \quad |b - \beta| \leq Cn\rho^3 + n^2\rho \|\eta\|_{L^2_{\text{loc}}}, \quad (3.44)$$

such that

$$\dot{q}_k = \sum_{l=1, \dots, N} D_{kl} |q_l|^2 q_k + iJ_k q_k + g_k, \quad (3.45)$$

$$\dot{\beta} = \sum_{1 \leq k \leq l \leq N} B_{kl} |z_k|^2 |z_l|^2 + g_b, \quad (3.46)$$

where $J_k(t)$ are real functions bounded by $C\rho^2$ to be defined in (3.93). The constants D_{kl} are complex and bounded by n^2 , with their real parts given by (3.72) satisfying

$$-5\gamma_0^+ n^2 \leq \text{Re } D_{kl} \leq Cn^4, \quad \text{Re } D_{kk} \leq -\gamma_0 n^2. \quad (3.47)$$

The order one constants γ_0 and γ_0^+ are defined in (1.5) and (3.2). The constants B_{kl} are real and $B_{kl} = -\frac{1}{2} c_1 \text{Re } D_{kl} + O(n^4)$. Moreover,

$$|g_k| \lesssim n\rho^4 + n^4\rho^3 + n\rho \|\eta^{(3)}\|_{L^2_{\text{loc}}} + \tilde{X}, \quad (3.48)$$

$$|g_b| \lesssim n\rho^5 + n^4\rho^4 + n\rho^2 \|\eta^{(3)}\|_{L^2_{\text{loc}}} + n^3\rho^2 \|\eta\|_{L^2_{\text{loc}}} + n^2 \|\eta\|_{L^2_{\text{loc}}}^2 + n \|\eta^3\|_{L^1_{\text{loc}}}, \quad (3.49)$$

where $\tilde{X} = \rho^2 \|\eta\|_{L^2_{\text{loc}}} + n \|\eta\|_{L^2_{\text{loc}}}^2 + \|\eta^3\|_{L^1_{\text{loc}}}$ is defined in (3.41).

PROOF: We first observe a consequence of Assumption A3: Suppose $k_0, k_1, \dots, k_j \in \{1, \dots, N\}$ with $j \leq j_{\max} = 3$ satisfy

$$\omega_{k_0} \pm \omega_{k_1} \pm \dots \pm \omega_{k_j} = 0. \quad (3.50)$$

Then j is odd and, after a relabeling, the two sets (with multiplicities) $\{k_0, \dots, k_{\frac{j-1}{2}}\}$ and $\{k_{\frac{j+1}{2}}, \dots, k_j\}$ are the same. The reason is as follows. Recall $\omega_k = e_k - e_0 + O(n^2) > 0$, and hence the expression in (3.50) is approximately equal to

$$(e_{k_0} - e_0) \pm (e_{k_1} - e_0) \pm \dots \pm (e_{k_j} - e_0). \quad (3.51)$$

For the expression in (3.50) to be zero, one of the ω_{k_m} must be with a negative sign. Hence two e_0 's in (3.51) cancel and (3.51) is the sum of j e_k 's minus the sum of another j e_k 's. Since $j \leq j_{\max}$, Assumption A3 then implies that they can be divided to two equal sets. Since there are $(j-1)$ e_0 's and all k_m 's are positive, j must be odd, these e_0 's cancel each other, and the other e_k 's form two equal sets.

Part 1. Excited states. Recall from (3.14) the equations for p_k :

$$\dot{p}_k = i^{-1} e^{i\omega_k t} \left[(u_k^+, F) + (u_k^-, \bar{F}) + [(u_k^+, h) + (u_k^-, \bar{h}) + (u_k, R)a] F_\theta \right], \quad (3.52)$$

where $h = \zeta_1 + \dots + \zeta_N + \eta$. The nonlinear terms F and F_θ are given in (3.15)–(3.16) and $F = F_1 + \dots + F_5$ is decomposed in (3.29). We shall derive the normal form of (3.52) using integration by parts for those terms no smaller than $n^2 \rho^3$.

Step 1 Integration of terms of order nz^2 .

Recall that the main part of F is $F_1 = \lambda Q(2\zeta\bar{\zeta} + \zeta^2)$. The main terms on the right side of (3.52) are among the first two groups of terms involving F ,

$$i^{-1} e^{i\omega_k t} \left\{ (u_k^+, F_1) + (u_k^-, \bar{F}_1) \right\}. \quad (3.53)$$

We denote the rest by R_k ,

$$\begin{aligned} R_k = i^{-1} e^{i\omega_k t} & \left[(u_k^+, F - F_1) + (u_k^-, \bar{F} - \bar{F}_1) \right. \\ & \left. + [(u_k^+, h) + (u_k^-, \bar{h}) + (u_k, R)a] F_\theta \right], \end{aligned} \quad (3.54)$$

and rewrite (3.52) as

$$\dot{p}_k = i^{-1} e^{i\omega_k t} \left\{ (u_k^+, F_1) + (u_k^-, \bar{F}_1) \right\} + R_k. \quad (3.55)$$

Substituting $F_1 = \sum_{l,m=1}^N \lambda Q(2\zeta_l \bar{\zeta}_m + \zeta_l \zeta_m)$ with $\zeta_l = z_l u_l^+ + \bar{z}_l u_l^-$ and using Convention 3.1, we get

$$\begin{aligned} & i^{-1} e^{i\omega_k t} \left\{ (u_k^+, F_1) + (u_k^-, \bar{F}_1) \right\} \\ &= \sum_{l,m=1}^N i^{-1} e^{i\omega_k t} \left\{ (\lambda Q u_k^+, 2\zeta_l \bar{\zeta}_m + \zeta_l \zeta_m) + (\lambda Q u_k^-, 2\zeta_l \bar{\zeta}_m + \bar{\zeta}_l \bar{\zeta}_m) \right\} \\ &= \sum_{l,m \in \Omega} c_k^{lm} e^{i\omega_k t} z_l \bar{z}_m, \end{aligned}$$

for some purely imaginary constants c_k^{lm} bounded by n . The phase factor of a typical term $e^{i\omega_k t} z_l z_m$ is $\omega_k - \omega_l - \omega_m$, which is nonzero by the observation for (3.50). Hence we can define

$$p_{k,1-} = \sum_{l,m \in \Omega} \frac{-ic_k^{lm}}{\omega_k - \omega_l - \omega_m} e^{i\omega_k t} z_l z_m, \quad (3.56)$$

$$p_{k,1e} = \sum_{l,m \in \Omega} \frac{ic_k^{lm}}{\omega_k - \omega_l - \omega_m} e^{i(\omega_k - \omega_l - \omega_m)t} (\dot{p}_l p_m + p_l \dot{p}_m), \quad (3.57)$$

and we have

$$\sum_{l,m \in \Omega} c_k^{lm} e^{i\omega_k t} z_l z_m = \dot{p}_{k,1-} + p_{k,1e}.$$

Because $p_{k,1e}$ are of order $n^2 z^3$, we need to extract their leading terms. Substitute (3.55) into (3.57) and collect terms. The leading terms in $p_{k,1e}$ are cubic polynomials in z_k and $p_{k,1e}$ can be rewritten as

$$p_{k,1e} = \sum_{l,m,n \in \Omega} i^{-1} e^{i\omega_k t} d_{k,1}^{lmj} z_l z_m z_j + g_{k,1}, \quad (3.58)$$

$$g_{k,1} = \sum_{l,m \in \Omega} \frac{ic_k^{lm}}{\omega_k - \omega_l - \omega_m} e^{i(\omega_k - \omega_l - \omega_m)t} (R_l p_m + p_l R_m). \quad (3.59)$$

Here $d_{k,1}^{lmj}$ are real constants bounded by n^2 , and the error terms $g_{k,1}$ are bounded by

$$|g_{k,1}| \leq C \max_{k,l,m} |c_k^{lm}| |p_l| |R_m| \leq C n \rho \max_m |R_m|. \quad (3.60)$$

We conclude

$$\dot{p}_k - \dot{p}_{k,1-} = R_k + \sum_{l,m,n \in \Omega} i^{-1} e^{i\omega_k t} d_{k,1}^{lmj} z_l z_m z_j + g_{k,1}. \quad (3.61)$$

Step 2 Integration of terms of the form $n z \eta^{(2)}$.

We now integrate $i^{-1} e^{i\omega_k t} (u_k^+, F_4)$, which is part of the first term of R_k . Recall $F_4 = 2\lambda Q[(\zeta + \bar{\zeta})\eta^{(2)} + \zeta \bar{\eta}^{(2)}]$ with $\eta^{(2)} = U^{-1} \sum_{l,m \in \Omega} z_l z_m \tilde{\eta}_{lm}$. Denote

$$\eta' = \sum_{l,m \in \Omega} z_l z_m \tilde{\eta}_{lm}, \quad (3.62)$$

where $\tilde{\eta}_{lm}$ are defined in (3.36) with $\|\tilde{\eta}_{lm}\|_{L_{\text{loc}}^2} \lesssim n$. Recall Lemma 2.5 that $U^{-1} = (U_+)^* - (U_-)^*$ **conj** and both U_+ and U_- commute with i and conjugation. Then

$\eta^{(2)} = U^{-1}\eta' = (U_+)^*\eta' - (U_-)^*\bar{\eta}'$ and

$$\begin{aligned} (u_k^+, F_4) &= (2\lambda Qu_k^+(\zeta + \bar{\zeta}), \eta^{(2)}) + (2\lambda Qu_k^+\bar{\zeta}, \bar{\eta}^{(2)}) \\ &= (2\lambda Qu_k^+(\zeta + \bar{\zeta}), [(U_+)^*\eta' - (U_-)^*\bar{\eta}']) \\ &\quad + (2\lambda Qu_k^+\bar{\zeta}, [((U_+)^*\eta' - (U_-)^*\bar{\eta}']) \\ &= (U_+[2\lambda Qu_k^+(\zeta + \bar{\zeta})] - U_-[2\lambda Qu_k^+\bar{\zeta}], \eta') \\ &\quad + (U_+[2\lambda Qu_k^+\bar{\zeta}] - U_-[2\lambda Qu_k^+(\zeta + \bar{\zeta})], \bar{\eta}'). \end{aligned} \quad (3.63)$$

$$(3.64)$$

Substituting $\zeta = \sum_j z_j u_j^+ + \bar{z}_j u_j^-$ and $\eta' = \sum_{l,m \in \Omega} z_l z_m \tilde{\eta}_{lm}$, we can write

$$i^{-1}e^{i\omega_k t}(u_k^+, F_4) = \sum_{l,m,j \in \Omega} d_k^{lmj} e^{i\omega_k t} z_l z_m z_j, \quad (3.65)$$

for some coefficients d_k^{lmj} bounded by n^2 . The phase factor of a typical term $e^{i\omega_k t} z_l z_m z_j$ is $\omega_k - \omega_l - \omega_m - \omega_j$. By the observation for (3.50), it is nonzero unless one of $|l|$, $|m|$, $|j|$ is k and the other two are the same. In this exceptional case $e^{i\omega_k t} z_l z_m z_j$ is of the form $e^{i\omega_k t} z_k z_l \bar{z}_l = |p_l|^2 p_k$. For fixed $l > 0$, there are six such terms if $l \neq k$, and three terms if $l = k$. We denote the sum of their coefficients as D_{kl} ,

$$\begin{aligned} D_{kl} &= d_k^{kl(-l)} + d_k^{k(-l)l} + d_k^{lk(-l)} + d_k^{(-l)kl} + d_k^{l(-l)k} + d_k^{(-l)lk}, \quad (l \neq k), \\ D_{kk} &= d_k^{kk(-k)} + d_k^{k(-k)k} + d_k^{(-k)kk}. \end{aligned} \quad (3.66)$$

The total of these zero-phase-factor terms is $\sum_{l=1}^N D_{kl} |p_l|^2 p_k$. The other terms can be integrated. Define

$$p_{k,2-} = \sum_{\omega_k - \omega_l - \omega_m - \omega_j \neq 0} \frac{-id_k^{lmj}}{\omega_k - \omega_l - \omega_m - \omega_j} e^{i\omega_k t} z_l z_m z_j, \quad (3.67)$$

$$g_{k,2} = - \sum_{\omega_k - \omega_l - \omega_m - \omega_j \neq 0} \frac{-id_k^{lmj}}{\omega_k - \omega_l - \omega_m - \omega_j} e^{i(\omega_k - \omega_l - \omega_m - \omega_j)t} \frac{d}{dt} (p_l p_m p_j). \quad (3.68)$$

We have

$$\sum_{l,m,j \in \Omega} d_k^{lmj} e^{i\omega_k t} z_l z_m z_j = \sum_{l=1}^N D_{kl} |z_l|^2 p_k + \frac{d}{dt} (p_{k,2-}) + g_{k,2}, \quad (3.69)$$

and

$$|g_{k,2}| \lesssim (\max |d_k^{lmj}|) \rho^2 \max_j |\dot{p}_j| \lesssim n^2 \rho^2 \max_j |\dot{p}_j|. \quad (3.70)$$

We now compute $\text{Re } D_{kl}$. We want to collect terms of the form $C e^{i\omega_k t} z_l \bar{z}_l z_k$ from (3.63)–(3.64) with $\text{Im } C \neq 0$. Hence we only need to consider those $\tilde{\eta}_{lm}$ with $\text{Im } \tilde{\eta}_{lm} \neq 0$, i.e., $l, m > 0$. The only terms from $\bar{\eta}'$ with resonance coefficients

are $\sum_{m,j>0} \overline{z_m z_j \tilde{\eta}_{mj}}$, which are of the form $\bar{z}_m \bar{z}_j$, with two bars. Hence the integral in (3.64) does not contain $z_l \bar{z}_l z_k$ and is irrelevant. From the integral in (3.63), we want to choose $z_l z_k$ from η' , i.e., $\sum_{m,j>0} z_m z_j \tilde{\eta}_{mj}$, and choose z_l from ζ or $\bar{\zeta}$ in $U_+[2\lambda Q u_k^+(\zeta + \bar{\zeta})] - U_-[2\lambda Q u_k^+ \bar{\zeta}]$. (Recall (2.1) that $(f, g) = \int \bar{f} g dx$.) For the second part we get $z_l \tilde{\Phi}_{kl}$ where

$$\tilde{\Phi}_{kl} = U_+ 2\lambda Q u_k^+(u_l^+ + u_l^-) - U_- 2\lambda Q u_k^+ u_l^-. \quad (3.71)$$

Since $\tilde{\eta}_{mj} = \tilde{\eta}_{jm}$ for $m, j > 0$, the terms with $z_l z_k$ in $\sum_{m,j>0} z_m z_j \tilde{\eta}_{mj}$ is $(2 - \delta_k^l) z_k z_l \tilde{\eta}_{kl}$. Therefore, also using the definition (3.36) of $\tilde{\eta}_{kl}$,

$$\begin{aligned} \operatorname{Re} D_{kl} &= \operatorname{Re} i^{-1} \int \tilde{\Phi}_{kl} (2 - \delta_k^l) \tilde{\eta}_{kl} dx \\ &= -(2 - \delta_k^l) \operatorname{Im} \left(\tilde{\Phi}_{kl}, \frac{1}{A - \omega_k - \omega_l - 0i} \mathbf{P}_c^A \Phi_{kl} \right). \end{aligned} \quad (3.72)$$

Recall

$$\begin{aligned} Q &= (1 + O(\varepsilon_0^2)) n \phi_0 + O(n^3), \quad u_k^+ = \phi_k + O(n^2), \quad u_k^- = O(n^2), \\ U_+ &= 1 + O(n^2), \quad U_- = O(n^2). \end{aligned} \quad (3.73)$$

Hence, by (3.35) and (3.71),

$$\Phi_{kl} = \lambda Q \phi_k \phi_l + O(n^3), \quad \tilde{\Phi}_{kl} = 2\lambda Q \phi_k \phi_l + O(n^3), \quad (3.74)$$

and we have

$$\operatorname{Re} D_{kl} = -2(2 - \delta_k^l)(1 + o(1))n^2 \operatorname{Im} \left(\phi_0 \phi_k \phi_l, \frac{1}{A - \omega_k - \omega_l - 0i} \mathbf{P}_c^A \phi_0 \phi_k \phi_l \right) + O(n^4). \quad (3.75)$$

By Assumption A2 and Lemma 2.5 (3), we have $-\operatorname{Re} D_{kl} \leq 5n^2 \gamma_0^+$ and $\operatorname{Re} D_{kl} \leq Cn^4$ for all $k, l > 0$, and $-\operatorname{Re} D_{kk} \geq \gamma_0 n^2$. We conclude (3.47).

Step 3 Integration of other terms.

We now integrate other terms in (3.61) no smaller than $n^2 z^3$. We first consider the first group of terms in R_k defined in (3.54). Using $F - F_1 = F_2 + F_3 + F_4 + F_5$ and removing $i^{-1} e^{i\omega_k t} (u_k^+, F_4)$, we get

$$i^{-1} e^{i\omega_k t} \left\{ (u_k^+, F_2 + F_3) + (u_k^-, \overline{F}_2) \right\} + g_{k,3},$$

where

$$g_{k,3} = i^{-1} e^{i\omega_k t} \left\{ (u_k^+, F_5) + (u_k^-, \overline{F}_3 + \overline{F}_4 + \overline{F}_5) \right\}. \quad (3.76)$$

We have, using $u_k^- = O(n^2)$ and the explicit form of $F_3 + F_4$,

$$|g_{k,3}| \lesssim \|F_5\|_{L_{\text{loc}}^1} + n^2 \|F_3 + F_4\|_{L_{\text{loc}}^1} \lesssim \|F_5\|_{L_{\text{loc}}^1} + n^2 (n^2 \rho^3). \quad (3.77)$$

We now consider the second group of terms in R_k , see (3.54),

$$i^{-1}e^{i\omega_k t} [(u_k^+, h) + (u_k^-, \bar{h}) + (u_k, R)a] F_\theta. \quad (3.78)$$

The only relevant terms are $i^{-1}e^{i\omega_k t} [(u_k^+, \zeta) + (u_k^-, \bar{\zeta})] F_\theta$. We move the other part to error term,

$$g_{k,4} = i^{-1}e^{i\omega_k t} [(u_k^+, \eta) + (u_k^-, \bar{\eta}) + (u_k, R)a] F_\theta. \quad (3.79)$$

Note that $(u_k, R) = O(n)$ since $u_k \perp Q$ and $R = Cn^{-2}Q + O(n)$ by Lemma 2.1. Also, $|(u_k^+, \eta) + (u_k^-, \bar{\eta})| \leq Cn^2 \|\eta\|_{L_{\text{loc}}^2}$ since $\text{Re } \eta \perp v_k$, $\text{Im } \eta \perp u_k$, and the differences between u_k, v_k, u_k^+, u_k^- are bounded by n^2 . Hence

$$|g_{k,4}| \leq (n^2 \|\eta\|_{L_{\text{loc}}^2} + n\rho^2) |F_\theta|. \quad (3.80)$$

Using $\zeta = \sum_l z_l u_l^+ + \bar{z}_l u_l^-$, we get

$$(u_k^+, \zeta) + (u_k^-, \bar{\zeta}) = \sum_{l=1}^N (C_k^l z_l + \tilde{C}_k^l \bar{z}_l),$$

where $C_k^l = (u_k^+, u_l^+) + (u_k^-, u_l^-)$ and $\tilde{C}_k^l = (u_k^-, u_l^+) + (u_k^+, u_l^-)$. Because $u_k^+ = \phi_k + O(n^2)$ and $u_k^- = O(n^2)$, see (3.73), we have

$$C_k^l = \delta_k^l + O(n^2), \quad \tilde{C}_k^l = O(n^2).$$

Thus, terms no smaller than $n^2 \rho^3$ in (3.78) are

$$i^{-1}e^{i\omega_k t} \left\{ z_k F_{\theta,1} + \sum_l [(C_k^l - \delta_k^l) z_l + \tilde{C}_k^l \bar{z}_l] F_{\theta,2} \right\}, \quad (3.81)$$

where $F_{\theta,1}$ and $F_{\theta,2}$ are leading parts of F_θ (3.16), (recall $a = b + a^{(2)}$)

$$F_{\theta,1} \equiv -[a + (c_1 R, \text{Re } F_1 + F_2)] \cdot [1 + (c_1 R, R)b]^{-1}, \quad (3.82)$$

$$F_{\theta,2} \equiv -[b + (c_1 R, \text{Re } F_1)], \quad (3.83)$$

and (3.78) is equal to $g_{k,4} + (3.81) + g_{k,5}$ where

$$g_{k,5} = i^{-1}e^{i\omega_k t} \left\{ z_k (F_\theta - F_{\theta,1}) + \sum_l [(C_k^l - \delta_k^l) z_l + \tilde{C}_k^l \bar{z}_l] (F_\theta - F_{\theta,2}) \right\}. \quad (3.84)$$

Since $|F_\theta - F_{\theta,2}| \leq |F_\theta - F_{\theta,1}| + |F_{\theta,1} - F_{\theta,2}|$, we have

$$|g_{k,5}| \lesssim \rho |F_\theta - F_{\theta,1}| + n^2 \rho |F_{\theta,1} - F_{\theta,2}|.$$

Recall (3.16) $F_\theta = -[a + (c_1 R, \text{Re } F)] \cdot [1 + (c_1 R, R)a + (c_1 R, \text{Re } h)]^{-1}$. Hence

$$\begin{aligned} F_\theta - F_{\theta,1} &= -(c_1 R, \text{Re } F_3 + F_4 + F_5) \cdot [1 + (c_1 R, R)b]^{-1} \\ &\quad + \frac{[a + (c_1 R, \text{Re } F)] \cdot [(c_1 R, R)a^{(2)} + (c_1 R, \text{Re } h)]}{[1 + (c_1 R, R)a + (c_1 R, \text{Re } h)] \cdot [1 + (c_1 R, R)b]}. \end{aligned} \quad (3.85)$$

Since $R = Cn^{-2}Q + O(n)$ and $\operatorname{Re} h \perp Q$, we have $|(c_1 R, \operatorname{Re} h)| \lesssim n \|h\|_{L^1_{\text{loc}}}$. Hence,

$$\begin{aligned} |F_\theta - F_{\theta,1}| &\lesssim n^{-1} \|F_3 + F_4 + F_5\|_{L^1_{\text{loc}}} + \left[|a| + n^{-1} \|F\|_{L^1_{\text{loc}}}\right] (\rho^2 + n \|h\|_{L^1_{\text{loc}}}) \\ &\lesssim n^{-1} (n^2 \rho^3 + X) + [\rho^2 + n^{-1} (n\rho^2 + X)] (n\rho + n \|\eta\|_{L^2_{\text{loc}}}) \\ &\lesssim n\rho^3 + n^{-1} X. \end{aligned}$$

Here we have used Lemma 3.2. We also have

$$\begin{aligned} F_{\theta,1} - F_{\theta,2} &= -[a^{(2)} + [c_1 R, \operatorname{Re} F_2]] \cdot [1 + (c_1 R, R)b]^{-1} \\ &\quad + [a + (c_1 R, \operatorname{Re} F_1 + F_2)] \cdot [1 + (c_1 R, R)b]^{-1} \cdot (c_1 R, R)b. \end{aligned} \quad (3.86)$$

Hence

$$|F_{\theta,1} - F_{\theta,2}| \lesssim (n^2 \rho^2 + n^{-1} \|F_2\|_{L^1_{\text{loc}}}) + (\rho^2 + n^{-1} \|F_1 + F_2\|_{L^1_{\text{loc}}}) n^{-2} \rho^2.$$

Since $\|F_1\|_{L^1_{\text{loc}}} \leq n\rho^2$ and $\|F_2\|_{L^1_{\text{loc}}} \leq \rho^3$, we have $|F_{\theta,1} - F_{\theta,2}| \lesssim n^2 \rho^2 + n^{-1} \rho^3$. Hence $g_{k,5}$ is bounded by

$$\begin{aligned} |g_{k,5}| &\lesssim \rho |F_\theta - F_{\theta,1}| + n^2 \rho |F_{\theta,1} - F_{\theta,2}| \\ &\lesssim \rho (n\rho^3 + n^{-1} X) + n^2 \rho (n^2 \rho^2 + n^{-1} \rho^3) \\ &\lesssim n\rho^4 + n^4 \rho^3 + n^{-1} \rho X. \end{aligned} \quad (3.87)$$

Summarizing, we can rewrite eq. (3.61) as

$$\frac{d}{dt} (p_k - p_{k,1-} - p_{k,2-}) = \sum_{l=1}^N D_{kl} |z_l|^2 p_k + \tilde{R}_k + \sum_{j=1}^5 g_{k,j}, \quad (3.88)$$

where

$$\begin{aligned} \tilde{R}_k &= i^{-1} e^{i\omega_k t} \left\{ \sum_{l,m,j \in \Omega} d_{k,1}^{lmj} z_l z_m z_j + (u_k^+, F_2 + F_3) + (u_k^-, \overline{F}_2) \right. \\ &\quad - z_k \frac{1}{1 + (c_1 R, R)b} [a + (c_1 R, \operatorname{Re} F_1 + F_2)] \\ &\quad \left. - \sum_l [(C_k^l - \delta_k^l) z_l + \tilde{C}_k^l \bar{z}_l] [b + (c_1 R, \operatorname{Re} F_1)] \right\}. \end{aligned} \quad (3.89)$$

We now integrate $\tilde{R}_k$. Denote

$$B = (c_1 R, R)b, \quad |B| \leq Cn^{-2} \rho^2. \quad (3.90)$$

Note that $\tilde{R}_k$ is of the form

$$ie^{i\omega_k t} [n^{-1} b^2 + bz + n^{-1} bz^2 + z^3 + n^{-1} z_k z^3], \quad (3.91)$$

in the sense that

$$\begin{aligned} \tilde{R}_k = \frac{ie^{i\omega_k t}}{1+B} & \left[f(B)n^{-1}b^2 + \sum_{j \in \Omega} f(B)bz_j + \sum_{j_1, j_2 \in \Omega} f(B)n^{-1}bz_{j_1}z_{j_2} \right. \\ & \left. + \sum_{j_1, j_2, j_3 \in \Omega} f(B)z_{j_1}z_{j_2}z_{j_3} + \sum_{j_1, j_2, j_3 \in \Omega} f(B)n^{-1}z_kz_{j_1}z_{j_2}z_{j_3} \right], \end{aligned} \quad (3.92)$$

where $f(B)$ are polynomials in B with real coefficients bounded by one. We have omitted their dependence on the summation indexes. The phase factors of the above summands are

$\omega_k, \quad \omega_k \pm \omega_{|j_1|}, \quad \omega_k \pm \omega_{|j_1|} \pm \omega_{|j_2|}, \quad \omega_k \pm \omega_{|j_1|} \pm \omega_{|j_2|} \pm \omega_{|j_3|}, \quad \pm \omega_{|j_1|} \pm \omega_{|j_2|} \pm \omega_{|j_3|},$
respectively. By the observation for (3.50), zero phase factor only occurs to bz and z^3 terms. Those terms with zero phase factor are

$$\frac{if_{bk}(B)}{1+B}e^{i\omega_k t}bz_k, \quad \frac{if_{kl}(B)}{1+B}e^{i\omega_k t}z_kz_l\bar{z}_l.$$

The sum of these terms is equal to $iJ_k e^{i\omega_k t} z_k = iJ_k p_k$, where

$$J_k(t) \equiv \frac{f_{bk}(B)}{1+B}b + \sum_{l=1}^N \frac{f_{kl}(B)}{1+B}|z_l|^2. \quad (3.93)$$

They are real functions bounded by $C\rho^2$. The other terms can be integrated. For example, if $j \in \Omega, j \neq k$, we have

$$\frac{if(B)}{1+B}e^{i\omega_k t}bz_j = \frac{d}{dt} \left(\frac{if(B)}{i(\omega_k - \omega_j)(1+B)}e^{i(\omega_k - \omega_j)t}bp_j \right) - \text{error},$$

where

$$\text{error} = e^{i(\omega_k - \omega_j)t} \frac{d}{dt} \left(\frac{if(B)}{i(\omega_k - \omega_j)(1+B)}bp_j \right),$$

and it is bounded by

$$|\text{error}| \lesssim |\dot{b}p_j| + |b\dot{p}_j| + |bp_j\dot{B}|.$$

Note $|\dot{B}| = Cn^{-2}|\dot{b}|$. We can integrate other terms similarly. Summing up, we get

$$\tilde{R}_k = iJ_k p_k + \frac{d}{dt} p_{k,3-} + g_{k,6}, \quad (3.94)$$

where $p_{k,3-}$ is of the same form (3.91) as $\tilde{R}_k$ and, adding the estimates for the integration remainders for all five kinds of terms in (3.91), we get

$$\begin{aligned} |g_{k,6}| & \lesssim n^{-1}(|\dot{b}\dot{b}| + b^2|\dot{B}|) + (|\dot{b}|\rho + \max_j |\dot{b}p_j| + |b\rho\dot{B}|) \\ & \quad + n^{-1}(|\dot{b}|\rho^2 + \max_j |b\rho\dot{p}_j| + |b\rho^2\dot{B}|) \\ & \quad + (\rho^2 \max_j |\dot{p}_j| + \rho^3|\dot{B}|) + n^{-1}(\rho^3 \max_j |\dot{p}_j| + \rho^4|\dot{B}|) \\ & \lesssim \rho^2 \max_j |\dot{p}_j| + \rho|\dot{b}|. \end{aligned} \quad (3.95)$$

Step 4 Final form. We now define

$$q_k \equiv p_k - p_{k,1-} - p_{k,2-} - p_{k,3-}. \quad (3.96)$$

Since $p_{k,1-} \sim nz^2$, $p_{k,2-} \sim n^2z^3$ and $p_{k,3-} \sim n^{-1}b^2 + bz + n^{-1}bz^2 + z^3 + n^{-1}z_kz^3$, we have

$$|p_k - q_k| \leq |p_{k,1-}| + |p_{k,2-}| + |p_{k,3-}| \leq Cn\rho^2. \quad (3.97)$$

From (3.89) and (3.94) we have

$$\begin{aligned} \dot{q}_k &= \sum_{l=1,\dots,N} D_{kl} |p_l|^2 p_k + iJ_k p_k + \sum_{j=1}^6 g_{k,j} \\ &= \sum_{l=1,\dots,N} D_{kl} |q_l|^2 q_k + iJ_k q_k + g_k, \end{aligned}$$

where

$$g_k = g_{k,1} + \dots + g_{k,6} + g_{k,7}, \quad (3.98)$$

$$g_{k,7} \equiv \sum_{l=1,\dots,N} D_{kl} (|p_l|^2 p_k - |q_l|^2 q_k) + iJ_k (p_k - q_k). \quad (3.99)$$

Since $D_{kl} = O(n^2)$ and $J_k = O(\rho^2)$, we have

$$|g_{k,7}| \lesssim n^2 \rho^2 (n\rho^2) + \rho^2 (n\rho^2) \lesssim n\rho^4. \quad (3.100)$$

Collecting estimates, we have

$$\begin{aligned} |g_k| &\lesssim \sum_{j=1}^7 |g_{k,j}| \lesssim n\rho \max_m |R_m| + n^2 \rho^2 \max_j |\dot{p}_j| + [\|F_5\|_{L_{\text{loc}}^1} + n^2(n^2 \rho^3)] \\ &\quad + (n^2 \|\eta\|_{L_{\text{loc}}^2} + n\rho^2) |F_\theta| + (n\rho^4 + n^4 \rho^3 + n^{-1} \rho X) \\ &\quad + (\rho^2 \max_j |\dot{p}_j| + \rho |\dot{b}|) + n\rho^4. \end{aligned}$$

Using $\max_m |R_m| \lesssim \|F - F_1\|_{L_{\text{loc}}^1} + (\rho + \|\eta\|_{L_{\text{loc}}^2}) |F_\theta|$ and Lemma 3.2, we have

$$\begin{aligned} |g_k| &\lesssim n\rho^4 + n^4 \rho^3 + n\rho \|F - F_1\|_{L_{\text{loc}}^1} + (n^2 \|\eta\|_{L_{\text{loc}}^2} + n\rho^2) |F_\theta| \\ &\quad + \|F_5\|_{L_{\text{loc}}^1} + n^{-1} \rho X \\ &\lesssim n\rho^4 + n^4 \rho^3 + n\rho(\rho^3 + X) + (n^2 \|\eta\|_{L_{\text{loc}}^2} + n\rho^2)(\rho^2 + n^{-1} X) \\ &\quad + (n\rho^4 + n\rho \|\eta^{(3)}\|_{L_{\text{loc}}^2} + \tilde{X}) + n^{-1} \rho X \\ &\lesssim n\rho^4 + n^4 \rho^3 + n\rho \|\eta^{(3)}\|_{L_{\text{loc}}^2} + \rho^2 \|\eta\|_{L_{\text{loc}}^2} + n \|\eta\|_{L_{\text{loc}}^2}^2 + \|\eta^3\|_{L_{\text{loc}}^1}. \end{aligned}$$

Part 2. Ground state.

We have derived the main oscillatory terms of $a(t)$ in (3.26)

$$a(t) = a^{(2)}(t) + b(t), \quad a^{(2)} = \sum_{k,l \in \Omega} a_{kl} z_k z_l,$$

with $b(t)$ given by (3.27). We have

$$\dot{b} = (c_1 Q, \text{Im}[F - F_1]) + (c_1 Q, \text{Im } h) F_\theta - A_{2,rm d}, \quad (3.101)$$

with $A_{2,rm d}$ given in (3.24),

$$A_{2,rm d} = \text{Im} \sum_{k,l=1}^N \left\{ e^{-i(\omega_k + \omega_l)s} i a_{kl,3} \frac{d}{ds} (p_k p_l) + e^{-i(\omega_k - \omega_l)s} i a_{kl,4} \frac{d}{ds} (p_k \bar{p}_l) \right\}.$$

As for the excited states p_k , we want to find a perturbation $\beta(t)$ of $b(t)$ so that oscillatory terms no smaller than $n^2 \rho^4$ on the right side of (3.101) are removed. We have observed in Lemma 3.2 that $|\dot{b}| \lesssim n \rho^3 + nX$ and $|(c_1 Q, \text{Im } h)| \lesssim n^3 \|h\|_{L^1_{\text{loc}}}$ since $h \in M$ is almost orthogonal to Q . Hence the right side of (3.101) is of the form

$$\begin{aligned} \dot{b} = & n \left\{ z^3 + b z + n^{-1} b z^2 + n^{-2} b^2 z + n^{-3} \eta + z^2 \eta^{(2)} + z^2 \eta^{(3)} + n \eta^2 + (\eta^3)_{\text{loc}} \right\} \\ & + n^3 z \left\{ z^2 + b + n^{-1} z (z^2 + b) + O(n^{-2} \rho^4) \right\} + n^3 \eta F_\theta \\ & + n^2 z \left\{ n z^2 + z^3 + b z + O(\rho^4/n) + n z \eta + z^2 \eta + n \eta^2 + (\eta^3)_{\text{loc}} + \dots \right\}. \end{aligned} \quad (3.102)$$

Here $(\eta^3)_{\text{loc}}$ means terms with same bound as $\|\eta^3\|_{L^1_{\text{loc}}}$. We shall calculate the normal form for b by integrating by parts those terms with orders which were crossed out. Notice that there are resonant terms with crossed-out orders, explicitly, terms of the form $n^2 |z_k|^2 |z_l|^2$. These terms cannot be integrated by parts and will remain on the right hand side. The final normal form equation is of the form (3.46). This procedure is the same as that for excited states and we shall not repeat it in details but point out a few key steps.

1. There are no terms of the form b^2 , b^3 or $|z_k|^2 b$ in the first line of (3.102). Terms of these forms are eliminated by the Im operator.

2. Terms of the form $C z_k z_l b^j$ are oscillatory if $k + l \neq 0$. These terms can be integrated. Similarly, by the observation for (3.50), terms of the form $C z_{m_1} z_{m_2} z_{m_3}$, $z_m b$ and $C z_{m_1} z_{m_2} z_{m_3} z_{m_4} z_{m_5}$ are also oscillatory with nonzero phase factors. These terms can be integrated.

3. Most terms of the form $C z_{m_1} z_{m_2} z_{m_3} z_{m_4}$ are oscillatory. The only terms with zero phase factor, $\omega_{m_1} + \dots + \omega_{m_4} = 0$, are of the form $C |z_k|^2 |z_l|^2$ by the observation for (3.50). These terms cannot be integrated and will remain in the final equation. Moreover, in order to survive the Im operator, these terms from $F - F_1$ must have complex coefficients, i.e., they must involve $\eta^{(2)}$ and are of the order $n z^2 \eta^{(2)}$.

4. We need to integrate terms of the form $n^2 z_j \eta$. They are from $(c_1 Q, \text{Im } 2\lambda Q[(\zeta + \bar{\zeta})\eta + \zeta \bar{\eta}])$ in the first line of (3.102). Since there is only one z_j involved, the Green's function is approximately $[H_0 - e_0 \pm (e_j - e_0)]^{-1}$ and is invertible in L^2 . Hence there is no resonance with the continuous spectrum. This integration is carried out in details in [19, p.193–195]. In contrast, resonant terms are of the form $n z_k z_l \eta^{(2)}$ and are mentioned in point 3.

In conclusion, we can find a perturbation $\beta(t)$ of $b(t)$ of the form

$$\begin{aligned} \beta = & b + \text{Re} \left\{ \sum C n z_{k_1} z_{k_2} z_{k_3} + C n z_k b + n^2 (z_k \psi_k, \eta) \right\} \\ & + \text{Re} \sum_{\omega_{k_1} + \dots + \omega_{k_4} \neq 0} C z_{k_1} z_{k_2} z_{k_3} z_{k_4} + \text{Re} \sum_{\omega_k + \omega_l \neq 0} C z_k z_l b \\ & + \text{Re} \left\{ C z_{k_1} \dots z_{k_5} / n + C z_{k_1} \dots z_{k_6} / n^2 \right\}, \end{aligned} \quad (3.103)$$

so that $\beta(t)$ satisfies a normal form equation,

$$\dot{\beta} = \sum_{1 \leq k \leq l \leq N} B_{kl} |z_k|^2 |z_l|^2 + g_b. \quad (3.104)$$

Here C denote complex constants bounded by one, and ψ_k denote some explicit local functions. B_{kl} are real constants bounded by n^2 , and g_b is an error term of the form

$$g_b \sim n z^5 + n^4 z^4 + n z^2 \eta^{(3)} + n^3 z^2 \eta + n^2 \eta^2 + n \eta^3 + \dots,$$

and we have the bounds $|\beta - b| \lesssim n \rho^3 + n^2 \rho \|\eta\|_{L_{\text{loc}}^2}$ and

$$|g_b| \lesssim n \rho^5 + n^4 \rho^4 + n \rho^2 \|\eta^{(3)}\|_{L_{\text{loc}}^2} + n^3 \rho^2 \|\eta\|_{L_{\text{loc}}^2} + n^2 \|\eta\|_{L_{\text{loc}}^2}^2 + n \|\eta^3\|_{L_{\text{loc}}^1}. \quad (3.105)$$

We now compute B_{kl} , the coefficients of $|z_k|^2 |z_l|^2$. The main contribution comes from

$$(c_1 Q, \text{Im } \lambda \zeta^2 \overline{\eta^{(2)}}), \quad (3.106)$$

where $\lambda \zeta^2 \overline{\eta^{(2)}}$ is from F_5 . Although there are terms from $(c_1 Q, \text{Im } 2\lambda \zeta \bar{\zeta} \eta^{(2)})$ and $(c_1 Q, \text{Im } 2\lambda Q[(\ell + \bar{\ell})\eta^{(2)} + \ell \bar{\eta}^{(2)}])$ with $\ell = a^{(2)} R + \eta^{(2)}$, their coefficients are small of order $O(n^4)$. Since $\eta^{(2)} = \sum_{k,l \in \Omega} U^{-1} z_k z_l \tilde{\eta}_{kl}$, and by (2.25) $U^{-1} = U_+^* - U_-^* \mathbf{conj}$, the expression is (3.106) is a sum of terms of the form $z_{k_1} z_{k_2} z_{k_3} z_{k_4}$. Moreover, since $\tilde{\eta}_{kl}$ has nontrivial imaginary part only if both $k, l > 0$, hence

$$\overline{\eta^{(2)}} = \sum_{k,l > 0} \mathbf{conj} U_+^* z_k z_l \tilde{\eta}_{kl} - \sum_{k,l > 0} U_-^* z_k z_l \tilde{\eta}_{kl} + \text{irrelevant terms}.$$

To get $|z_k|^2 |z_l|^2$ from $\text{Im } \lambda \zeta^2 \overline{\eta^{(2)}}$, the relevant terms in ζ^2 are $\nu_{kl}(z_k u_k^+ z_l u_l^+)$ and $\nu_{kl}(\bar{z}_k u_k^- \bar{z}_l u_l^-)$, where $\nu_{kl} \equiv 2 - \delta_k^l$. Therefore

$$\begin{aligned} (c_1 Q, \text{Im } \lambda \zeta^2 \overline{\eta^{(2)}}) &= \sum_{k,l > 0} (c_1 Q, \text{Im } \lambda \nu_{kl}(z_k u_k^+ z_l u_l^+) \bar{z}_k \bar{z}_l \mathbf{conj} U_+^* \tilde{\eta}_{kl}) \\ &+ \sum_{k,l > 0} (c_1 Q, \text{Im } \lambda \nu_{kl}(\bar{z}_k u_k^- \bar{z}_l u_l^-) z_k z_l (-1) U_-^* \tilde{\eta}_{kl}) + (*), \end{aligned}$$

where $(*)$ denotes terms of the form $z_{k_1} z_{k_2} z_{k_3} z_{k_4}$ with $\omega_{k_1} + \dots + \omega_{k_4} \neq 0$. Irrelevant terms with $\omega_{k_1} + \dots + \omega_{k_4} = 0$ are eliminated by the Im operator. Moreover, we can disregard the second sum since $U_- = O(n^2)$ is smaller than U_+ . By (3.73) and $\tilde{\eta}_{kl} = O(n)$, we have

$$\begin{aligned} B_{kl} &= (U_+ (c_1 Q \lambda \nu_{kl} u_k^+ u_l^+) , -\text{Im } \tilde{\eta}_{kl}) + O(n^4) \\ &= c_1 \nu_{kl} \left(U_+ (\lambda Q \phi_k \phi_l) , \text{Im } \frac{1}{A - \omega_k - \omega_l - 0i} \mathbf{P}_c^A \Phi_{kl} \right) + O(n^4). \end{aligned}$$

In view of (3.72)–(3.75), we have

$$B_{kl} = -\frac{c_1}{2} \text{Re } D_{kl} + O(n^4). \quad (3.107)$$

The proof of Lemma 3.3 is complete. **Q.E.D.**

3.4 Main estimates

Theorem 3.1 can be proved using the following proposition and a continuity argument. Recall $\rho(t) \equiv [\rho_0^{-2} + N^{-1} \gamma_0 n^2 t]^{-1/2}$, $\Lambda(t) = (1+s)^{-1/2} \rho^2(s)$, and $D = 6N|c_1| \gamma_0^+ / \gamma_0$.

Proposition 3.4 *Suppose the assumptions of Theorem 3.1 hold. Suppose for a fixed $T > 0$ we can find the best approximation $Q_{E(T)}$ of $\psi(T)$, i.e., $a_E(T) = 0$ in the decomposition (3.1) of $\psi(T)$ with $E = E(T)$. Define*

$$\begin{aligned} M_T &\equiv \sup_{0 \leq t \leq T} \max \left\{ \rho(t)^{-1} (\sum_{k=1}^N |z_k(t)|^2)^{1/2}, \quad 2D^{-1} \rho^{-2}(t) |a(t)|, \right. \\ &\quad n^{-4/5} \rho(t)^{-8/5} \|\eta(t)\|_{L^5}, \quad [\Lambda(t) + n^{1/3} \rho^{8/3}(t)]^{-1} \|\eta^{(3)}(t)\|_{L_{\text{loc}}^2}, \\ &\quad \left. 2(3D)^{-1} \rho_0^{-2} |E(T) - E_0| \right\}. \end{aligned} \quad (3.108)$$

Suppose, furthermore, $M_T \leq 2$. Then we have $M_T \leq 3/2$.

PROOF: Since $M_T \leq 2$, we have $|E(T) - E_0| \leq 3D\rho_0^2$ and, for $t \in [0, T]$,

$$\begin{aligned} (\sum_{k=1}^N |z_k(t)|^2)^{1/2} &\leq 2\rho(t), \quad |a(t)| \leq D\rho(t)^2, \\ \|\eta(t)\|_{L^5} &\leq 2n^{4/5} \rho(t)^{8/5}, \\ \|\eta^{(3)}(t)\|_{L_{\text{loc}}^2} &\leq 2\Lambda(t) + 2n^{1/3} \rho^{8/3}(t). \end{aligned} \quad (3.109)$$

Since $\eta = \eta^{(2)} + \eta^{(3)}$, $\|\eta^{(2)}\|_{L_{\text{loc}}^2} \lesssim n\rho^2$ and $\rho \leq \varepsilon_0 n$,

$$\|\eta(t)\|_{L_{\text{loc}}^2} \leq Cn\rho^2 + \|\eta^{(3)}(t)\|_{L_{\text{loc}}^2} \leq Cn\rho^2 + 2\Lambda(t). \quad (3.110)$$

Note that $\Lambda(t) \leq \rho^2$ by its definition. Therefore $\|\eta(t)\|_{L_{\text{loc}}^2} \leq C\rho^2$, $\|\eta(t)\|_{L_{\text{loc}}^2 \cap L^5} \leq Cn^{2/5} \rho^{8/5}$, and the assumptions of Lemmas 3.2 and 3.3 are satisfied with $\rho = \rho(t)$.

We have, using Lemma 2.2, (3.4) and (3.109),

$$|E(T) - E_0| \leq |E(T) - E(0)| + |E(0) - E_0| \leq \frac{9}{8}[|a_{E(T)}(0)| + |a_{E_0}|] \leq \frac{9}{4}D\rho_0^2. \quad (3.111)$$

Since $\|\psi(t)\|_{L^2} = \|\psi_0\|_{L^2} \ll 1$, we have $\|\eta(t)\|_{L^2} \ll 1$. By Hölder inequality,

$$\begin{aligned} \|\eta^3\|_{L^{5/4}} &\lesssim \|\eta\|_{L^2}^{2/3} \|\eta\|_{L^5}^{7/3} \lesssim o(1)(n^{4/5}\rho(t)^{8/5})^{7/3} = o(1)n^{28/15}\rho^{56/15}, \\ \|\eta^3\|_{L^1} &\lesssim \|\eta\|_{L^2}^{4/3} \|\eta\|_{L^5}^{5/3} \lesssim o(1)(n^{4/5}\rho(t)^{8/5})^{5/3} = o(1)n^{4/3}\rho^{8/3}, \\ \|\eta^3\|_{L_{\text{loc}}^1} &\lesssim \|\eta\|_{L_{\text{loc}}^2}^{4/3} \|\eta\|_{L^5}^{5/3} \lesssim \|\eta\|_{L_{\text{loc}}^2}^{4/3} n^{4/3}\rho^{8/3}. \end{aligned} \quad (3.112)$$

Recall $\tilde{X} = \rho^2 \|\eta\|_{L_{\text{loc}}^2}^2 + n \|\eta\|_{L_{\text{loc}}^2}^2 + \|\eta^3\|_{L_{\text{loc}}^1}$ and $X \lesssim n\rho \|\eta\|_{L_{\text{loc}}^2} + \tilde{X}$. We have

$$\rho^2 \|\eta\|_{L_{\text{loc}}^2}^2 + n \|\eta\|_{L_{\text{loc}}^2}^2 \lesssim \rho^2(n\rho^2 + \Lambda) + n(n\rho^2 + \Lambda)^2 \lesssim n\rho^4 + \rho^2\Lambda + n\Lambda^2. \quad (3.113)$$

Since $\|\eta^3\|_{L_{\text{loc}}^1} \ll \rho^2 \|\eta\|_{L_{\text{loc}}^2}^2$ by (3.112), we have

$$\tilde{X} \lesssim n\rho^4 + \rho^2\Lambda + n\Lambda^2, \quad X \lesssim n^2\rho^3 + n\rho\Lambda + n\Lambda^2. \quad (3.114)$$

We now estimate $\eta(t)$. Recall from (3.33), (3.31) that $\eta(t) = U^{-1}e^{-i\theta(t)}\tilde{\eta}(t)$ and

$$\tilde{\eta}(t) = Ue^{t\mathcal{L}}\eta(0) + \int_0^t e^{-iA(t-s)} \mathbf{P}_c^A F_\eta(s) ds.$$

By Lemma 3.2, (3.109), (3.112), $\|\eta(t)\|_{L_{\text{loc}}^2} \leq C\rho^2$, and $\|\eta(t)\|_{L_{\text{loc}}^2 \cap L^5} \leq Cn^{2/5}\rho^{8/5}$,

$$\begin{aligned} \|F_\eta\|_{L^{5/4}} &\lesssim n\rho^2 + n\rho \|\eta\|_{L_{\text{loc}}^2} + n \|\eta\|_{L_{\text{loc}}^2 \cap L^5}^2 + \|\eta^3\|_{L^{5/4}} + n^2\rho^2 \|\eta\|_{L^5} \\ &\lesssim n\rho^2 + n\rho(\rho^2) + n(n^{2/5}\rho^{8/5})^2 + n^{28/15}\rho^{56/15} + n^2\rho^2(n^{4/5}\rho^{8/5}) \lesssim n\rho^2. \end{aligned}$$

Using (3.5) for $\eta(0)$, the decay estimates of $e^{t\mathcal{L}}$ and e^{-itA} , and the boundedness in Sobolev spaces of U and U^{-1} from Lemma 2.5, we have

$$\begin{aligned} \|\eta(t)\|_{L^5} &\leq \|e^{t\mathcal{L}}\eta(0)\|_{L^5} + C \int_0^t |t-s|^{-9/10} \|F_\eta(s)\|_{L^{5/4}} ds \\ &\leq n^{4/5}\rho^{8/5} + C \int_0^t |t-s|^{-9/10} n\rho(s)^2 ds \leq n^{4/5}\rho(t)^{8/5} + Cn^{4/5}\rho_0^{2r}\rho^{9/5-2r}, \end{aligned}$$

for any $r > 0$. Here we have used $\rho(s) = Cn^{-1}(\Delta t + s)^{-1/2}$ with $\Delta t = Cn^{-2}\rho_0^{-2}$ and

$$\int_0^t |t-s|^{-9/10} (\Delta t + s)^{-1} ds \leq \Delta t^{-r} (\Delta t + t)^{-9/10+r}, \quad (3.115)$$

for any $r > 0$. Taking $2r = 1/5$, we get $\|\eta(t)\|_{L^5} \leq \frac{3}{2}n^{4/5}\rho(t)^{8/5}$.

We now consider the L_{loc}^2 estimates of η . Recall $\eta^{(3)} = \eta_1^{(3)} + \dots + \eta_4^{(3)}$. The estimate of $\eta_1^{(3)}$ is by (3.5) and that $U^{-1}e^{-i\theta}U = e^{-i\theta} + O(1)[i, U] = e^{-i\theta} + O(n^2)$,

$$\left\| \eta_1^{(3)}(t) \right\|_{L_{\text{loc}}^2} \leq (1 + o(1))\Lambda(t).$$

By the singular decay estimate (2.30) in Lemma 2.5 and the definition (3.36) of $\tilde{\eta}_{kl}$ with $\tilde{\eta}_{kl} = O(n)$, we have

$$\left\| \eta_2^{(3)}(t) \right\|_{L_{\text{loc}}^2} \lesssim n\rho_0^2(1+t)^{-3/2} \ll \Lambda(t),$$

$$\left\| \eta_3^{(3)}(t) \right\|_{L_{\text{loc}}^2} \lesssim \int_0^t \langle t-s \rangle^{-3/2} n(\rho^2|\dot{\theta}| + \rho \max_k |\dot{p}_k|) ds.$$

By the bounds of $\dot{\theta}$ and $\max_j |\dot{p}_j|$ in Lemma 3.2, and $\Lambda \leq \rho^2$, we have

$$\begin{aligned} n(\rho^2|\dot{\theta}| + \rho \max_k |\dot{p}_k|) &\lesssim n\rho^2(\rho^2 + n^{-1}X) + n\rho(n\rho^2 + X) \lesssim n^2\rho^3 + n\rho X \\ &\lesssim n^2\rho^3 + n\rho(n^2\rho^3 + n\rho\Lambda + n\Lambda^2) \lesssim n^2\rho^3. \end{aligned}$$

Hence

$$\left\| \eta_3^{(3)}(t) \right\|_{L_{\text{loc}}^2} \lesssim \int_0^t \langle t-s \rangle^{-3/2} n^2\rho^3(s) ds \lesssim n^2\rho^3(t).$$

Finally, by Lemma 3.2, (3.112), $\|\eta(t)\|_{L_{\text{loc}}^2 \cap L^5} \leq Cn^{2/5}\rho^{8/5}$, and $\Lambda \leq \rho^2$,

$$\begin{aligned} \|F_{\eta,3}\|_{L^{5/4} \cap L^1} &\lesssim \rho^3 + n\rho \|\eta\|_{L_{\text{loc}}^2} + n \|\eta\|_{L_{\text{loc}}^2 \cap L^5}^2 + \|\eta^3\|_{L^{5/4} \cap L^1} + n^2\rho^2 \|\eta\|_{L^5} \\ &\lesssim \rho^3 + n\rho(n\rho^2 + \Lambda) + n(n^{2/5}\rho^{8/5})^2 + o(1)n^{4/3}\rho^{8/3} + n^2\rho^2(n^{4/5}\rho^{8/5}) \\ &\lesssim \rho^3 + n^{4/3}\rho^{8/3} + n\rho\Lambda. \end{aligned}$$

Hence, bounding the integrand of $\eta_4^{(3)}$ by either L^∞ or L^5 -norm, we have

$$\begin{aligned} \left\| \eta_4^{(3)}(t) \right\|_{L_{\text{loc}}^2} &\leq \int_0^t \min \{ |t-s|^{-3/2}, |t-s|^{-9/10} \} \|F_{\eta,3}(s)\|_{L^{5/4} \cap L^1} ds \\ &\lesssim \int_0^t \min \{ |t-s|^{-3/2}, |t-s|^{-9/10} \} [\rho^3 + n^{4/3}\rho^{8/3} + n\rho\Lambda](s) ds \\ &\lesssim [\rho^3 + n^{4/3}\rho^{8/3} + n\rho\Lambda](t). \end{aligned}$$

Summing the estimates, we conclude

$$\left\| \eta^{(3)}(t) \right\|_{L_{\text{loc}}^2} \leq \sum_{j=1}^4 \left\| \eta_j^{(3)} \right\|_{L_{\text{loc}}^2} \leq \frac{5}{4}\Lambda + C\rho^3 + Cn^{4/3}\rho^{8/3} \leq \frac{5}{4}(\Lambda(t) + n^{1/3}\rho^{8/3})(t).$$

We now estimate the error terms g_k and g_b . Using $\rho \leq \varepsilon_0 n$, (3.48), (3.49), (3.109)–(3.114) and $\Lambda \leq \rho^2$, we have

$$\begin{aligned} |g_k| &\lesssim n\rho^4 + n^4\rho^3 + n\rho \left\| \eta^{(3)} \right\|_{L_{\text{loc}}^2} + \tilde{X} \\ &\lesssim n\rho^4 + n^4\rho^3 + n\rho(\Lambda + n^{1/3}\rho^{8/3}) + (n\rho^4 + \rho^2\Lambda + n\Lambda^2) \\ &\lesssim (\varepsilon_0^{2/3} + n^2)n^2\rho^3 + n\rho\Lambda. \end{aligned} \tag{3.116}$$

$$\begin{aligned}
|g_b| &\lesssim n\rho^5 + n^4\rho^4 + n\rho^2 \|\eta^{(3)}\|_{L_{\text{loc}}^2} + n^3\rho^2 \|\eta\|_{L_{\text{loc}}^2} + n^2 \|\eta\|_{L_{\text{loc}}^2}^2 + n \|\eta^3\|_{L_{\text{loc}}^1} \\
&\lesssim n\rho^5 + n^4\rho^4 + n\rho^2(\Lambda + n^{4/3}\rho^{8/3}) + n^3\rho^2(n\rho^2 + \Lambda) + n^2(n\rho^2 + \Lambda)^2 \\
&\quad + n(n\rho^2 + \Lambda)^{4/3}n^{4/3}\rho^{8/3} \\
&\lesssim (\varepsilon_0^{2/3} + n^2)n^2\rho^4 + n\rho^2\Lambda.
\end{aligned} \tag{3.117}$$

We now estimate $a(t)$. By (3.44) of Lemma 3.3 we have

$$|\beta - a| \leq |\beta - b| + |b - a| \leq (Cn\rho^3 + Cn^2\rho \|\eta\|_{L_{\text{loc}}^2}) + Cn^2\rho^2.$$

Since $\|\eta\|_{L_{\text{loc}}^2} \lesssim \rho^2$, we get $|\beta - a| \leq Cn^2\rho^2$. Since $a(T) = 0$, we have $|\beta(T)| = |\beta(T) - a(T)| \leq Cn^2\rho^2(T)$. Using (3.46) and (3.117) for g_b , we have

$$\begin{aligned}
|\beta(t)| &\leq |\beta(T)| + \int_t^T \sum |B_{kl}| |z_k|^2 |z_l|^2 + |g_b(s)| ds \\
&\leq Cn^2\rho^2(T) + \int_t^T (\max_{k,l} |B_{kl}|) \rho^4 + C \left[(\varepsilon_0^{2/3} + n^2)n^2\rho^4 + n\rho^2\Lambda \right] (s) ds \\
&\leq \frac{1}{2}D\rho^2(t) + o(1)\rho^2(t) \leq \frac{5}{8}D\rho^2(t).
\end{aligned}$$

Here we have used $\rho(s) = (\rho_0^{-2} + N^{-1}\gamma_0 n^2 s)^{-1/2}$ and $(\max_{k,l} |B_{kl}|)/(N^{-1}\gamma_0 n^2) \leq D/2$. We also have used $\Lambda(s) = n^{1/2}(1+s)^{-1/2}\rho^2(s)$, $\rho(s) = Cn^{-1}(\Delta t + s)^{-1/2}$ with $\Delta t = C(n\rho_0)^{-2}$, and the following estimate: for $t \geq 0$ and $\Delta t \geq 1$, $m, r > 0$, $m + r > 1$,

$$\int_t^\infty (\Delta t + s)^{-m}(1+s)^{-r} ds \leq \int_t^\infty (\Delta t + s)^{-m-r} ds \leq C(\Delta t + s)^{-m-r+1}. \tag{3.118}$$

Hence

$$|a(t)| \leq |\beta(t)| + |a(t) - \beta(t)| \leq \frac{5}{8}D\rho^2(t) + Cn^2\rho^2(t) \leq \frac{3}{4}D\rho^2(t).$$

We now estimate the excited states $z_k(t)$. For their initial value, we have

$$\|(\zeta_E - \zeta_{E_0})(0)\| \leq Cn^{-1}(\rho_0/n)|E - E_0| \leq Cn^{-2}\rho_0^3, \tag{3.119}$$

by (2.3) of Lemma 2.2 and $|E - E_0| \leq 3D\rho_0^2$. In particular, $(\sum_{k=1}^N z_{k,E}(0)^2)^{1/2} \leq \frac{5}{8}\rho_0$ by (3.4). Let $f_k = |q_k|^2$. By (3.44), $f(0) \leq \frac{1}{2}\rho_0^2$. By (3.45) and that $J_k(t)$ are real, we have

$$\dot{f}_k = \sum_{l=1}^N 2 \operatorname{Re} D_{kl} f_l f_k + 2 \operatorname{Re} \bar{q}_k g_k, \quad (k = 1, \dots, N). \tag{3.120}$$

Let $f = f_1 + \dots + f_N$. Since $|\operatorname{Re} D_{kl}| \leq 5\gamma_0^+ n^2$, summing (3.120) over k we get

$$|\dot{f}| \leq (10\gamma_0^+ n^2)f^2 + 2N \max_k |q_k g_k| \leq Cn^2\rho^4 + C\rho((\varepsilon_0^{2/3} + n^2)n^2\rho^3 + n\rho\Lambda),$$

by (3.116). Let $t_0 = n^{-3} \ll n^{-2}\rho_0^{-2}$. For $t \in [0, t_0]$, $\Lambda(t) \leq \rho_0^2(1+t)^{-1/2}$ and

$$|f(0) - f(t)| \lesssim \int_0^t n^2 \rho^4 + n \rho^2 \Lambda ds \lesssim n^2 \rho_0^4 t_0 + n \rho_0^4 \sqrt{t_0} \ll \rho_0^2,$$

since $t_0 \ll n^{-2}\rho_0^{-2}$. We conclude $f(t) \sim f(0)$ for $t \in [0, t_0]$. For $t \geq t_0$, we have

$$|g_k(t)| \lesssim (\varepsilon_0^{2/3} + n^2)n^2 \rho^3 + n \rho \Lambda \lesssim (\varepsilon_0^{2/3} + n^{1/2})n^2 \rho^3. \quad (3.121)$$

Since the positive part of $\text{Re } D_{kl}$ is bounded by n^4 and $\text{Re } D_{kk} \leq -\gamma_0 n^2$, we have

$$\dot{f} \leq -\sum_{k=1}^N 2\gamma_0 n^2 f_k^2 + Cn^4 f^2 + 2N \max_k |q_k g_k| \leq -\frac{3}{2N} \gamma_0 n^2 f^2 + 2N \max_k |q_k g_k|.$$

Using $\rho^2(t) = (\rho_0^{-2} + N^{-1}\gamma_0 n^2 t)^{-1}$ and (3.121), we have

$$\frac{d}{dt}(\rho^2) = -N^{-1}\gamma_0 n^2 \rho^4 \geq -\frac{3}{2N} \gamma_0 n^2 (\rho^2)^2 + 2N \max_k |q_k g_k|.$$

Since we have shown previously that $\rho^2(t) \geq f(t)$ for $t \in [0, t_0]$, in particular $\rho^2(t_0) \geq f(t_0)$, we have $\rho^2(t) \geq f(t)$ for all $t \geq t_0$ by comparison principle. Thus we have $\rho^2(t) \geq f(t)$ for all $t \geq 0$. **Q.E.D.**

We now prove Theorem 3.1 using Proposition 3.4 and a continuity argument.

PROOF OF THEOREM 3.1: By the assumption of Theorem 3.1, we have $M_T \leq 3/2$ for $T = 0$. Denote by J the set of all T such that we can find a best approximation $Q_{E(T)}$ of $\psi(T)$ and we have $M_T \leq 3/2$ with respect to $E = E(T)$. Clearly J is a closed interval containing 0 by the continuity of the Schrödinger equations.

Suppose $T' \in J$. By continuity there is a $\delta > 0$ which may depend on ψ_0 and T' such that, for all $T \in [T', T' + \delta]$, there is a best approximation $Q_{E(T)}$ of $\psi(T)$, and with respect to $E = E(T)$ we have $M_T \leq 2$. By Proposition 3.4, we have $M_T \leq 3/2$. Hence $(0, T' + \delta) \subset J$. This shows the right end of J is open and hence $J = [0, \infty)$.

For $t < T$, we have $|E(t) - E(T)| \leq \frac{5}{4}|a_T(t)| \leq D\rho(t)^2$ by Lemma 2.2 and Proposition 3.4. This uniform bound shows $E(t)$ has a unique limit E_∞ as $t \rightarrow \infty$ and $|E(t) - E_\infty| \leq D\rho(t)^2$. By continuity of M_T in T we have $M_\infty \leq 3/2$.

Finally we prove the lower bound (3.8). If $(\sum_{k=1}^N \|\zeta_k\|_{L^2}^2)^{1/2} \geq \frac{1}{4}\rho_0$, we have $f(0) \geq C\rho_0^2$. We have shown previously that $f(t) \sim f(0)$ for $t \leq t_0$. Summing (3.120) over k and using the error estimates, we have, for $t \geq t_0$,

$$\dot{f} \geq -10\gamma_0^+ n^2 f^2 + 2N \max_k |q_k g_k| \geq -12\gamma_0^+ n^2 f^2.$$

By a similar comparison argument we have $f(t) \geq (f(t_0)^{-1} + 12\gamma_0^+ n^2 (t - t_0))^{-1}$. Hence $f(t) \geq C\rho^2(t)$ for all $t \geq 0$. This completes the proof of Theorem 3.1. **Q.E.D.**

We remark that, since B_{kl} are (almost) positive, $\beta(t)$ and hence $a(t)$ are increasing as $t \rightarrow \infty$. This shows that the ground state gains energy from excited states even in the stabilization regime.

4 Transition Regimes

In this section we study the dynamics in the transition regime. We will prove Theorem 1.1 using Theorem 3.1. In this regime the natural operator is $H_0 = -\Delta + V$ and the natural decomposition is

$$\psi = x_0\phi_0 + \cdots + x_N\phi_N + \xi, \quad \xi \in \mathbf{H}_c(H_0).$$

4.1 Equations

Recall $H_0 = -\Delta + V$ has $N + 1$ simple eigenvalues e_k , $k = 0, \dots, N$, with corresponding normalized eigenvectors ϕ_k : $H_0\phi_k = e_k\phi_k$, $\|\phi_k\|_2 = 1$. We can decompose the solutions $\psi(t) = \psi(t, x)$ respect to H_0 :

$$\psi(t) = \chi(t) + \xi(t), \quad \chi(t) = \sum_{k=0}^N x_k(t)\phi_k, \quad (4.1)$$

where $x_k(t) \in \mathbb{C}$ and $\xi(t) \in \mathbf{H}_c(H_0)$. Substituting (4.1) into (1.1), we obtain the following system for these components:

$$\begin{cases} i\dot{x}_k = e_k x_k + (\phi_k, G), & (k = 0, \dots, N), \\ i\partial_t \xi = H_0 \xi + \mathbf{P}_c G, & G = \lambda \psi^2 \bar{\psi}, \end{cases} \quad (4.2)$$

with initial conditions $x_k(0) = x_k^0$ and $\xi(0) = \xi_0$. From these equations we find that each $x_k(t)$ is oscillatory with a main oscillation factor $e^{-ie_k t}$. We say that the *phase factor* of x_k is $-e_k$. Define $u_k(t)$ by

$$x_k(t) = e^{-ie_k t} u_k(t). \quad (4.3)$$

The function $u_k(t)$ has the same magnitude as $x_k(t)$ but is not as oscillatory. In particular, $|\dot{u}_k|$ is smaller than $|\dot{x}_k|$. We shall study the following system for u_k and ξ , which is equivalent to (4.2):

$$\dot{u}_k(t) = e^{ie_k t} (\phi_k, G(t)), \quad (k = 0, \dots, N), \quad (4.4)$$

$$\xi(t) = e^{-iH_0 t} \xi_0 + \int_0^t e^{-iH_0(t-s)} \mathbf{P}_c i^{-1} G(s) ds, \quad G = \lambda \psi^2 \bar{\psi}. \quad (4.5)$$

4.2 Decompositions of G and ξ

It is useful to decompose various terms according to their orders in n , so that we can identify the main terms. We expect that $x_k = O(n)$ and $\xi = O(n^3)$ locally after an

initial layer of time. We first decompose $G = \psi^2 \bar{\psi}$. Using (4.1) that $\psi = \chi + \xi$ with $\chi = \sum_{k=0}^N x_k \phi_k$, we decompose G as

$$G = \lambda \psi^2 \bar{\psi} = \lambda (\chi + \xi)^2 (\bar{\chi} + \bar{\xi}) = G_3 + G_5 + G_7, \quad (4.6)$$

where

$$G_3 = \lambda \chi^2 \bar{\chi}, \quad (4.7)$$

$$G_5 = \lambda \chi^2 \bar{\xi} + 2\lambda |\chi|^2 \xi, \quad (4.8)$$

$$G_7 = \lambda \bar{\chi} \xi^2 + 2\lambda \chi |\xi|^2 + \lambda \xi^2 \bar{\xi}. \quad (4.9)$$

Note that $G_3 = O(n^3)$, $G_5 = O(n^5)$ and $G_7 = O(n^7)$.

We now identify the main term of ξ using the integral equation (4.5). The main term of the integrand is $i^{-1}G_3$. Using (4.3) and factoring out the main phase factors in G_3 we have

$$G_3(t) = \sum_{l,m,j=0}^N e^{i(-e_l - e_m + e_j)t} u_l u_m \bar{u}_j(t) \phi_{lmj}, \quad (4.10)$$

where

$$\phi_{lmj} = \lambda \phi_l \phi_m \phi_j. \quad (4.11)$$

We now integrate by parts a typical term in $i^{-1}G_3$:

$$\begin{aligned} & \int_0^t e^{-iH_0(t-s)} \mathbf{P}_c i^{-1} e^{i(-e_l - e_m + e_j)s} u_l u_m \bar{u}_j(s) \phi_{lmj} ds \\ &= \lim_{r \rightarrow 0+} i^{-1} e^{-iH_0 t} \int_0^t e^{is(H_0 - e_l - e_m + e_j - ri)} u_l u_m \bar{u}_j(s) \mathbf{P}_c \phi_{lmj} ds \\ &= \lim_{r \rightarrow 0+} i^{-1} e^{-iH_0 t} \frac{1}{i(H_0 - e_l - e_m + e_j - ri)} \\ & \quad \left\{ \left[e^{iH_0 s} e^{i(-e_l - e_m + e_j)s} u_l u_m \bar{u}_j(s) \mathbf{P}_c \phi_{lmj} \right]_0^t \right. \\ & \quad \left. - \int_0^t e^{iH_0 s} e^{i(-e_l - e_m + e_j)s} \frac{d}{ds} (u_l u_m \bar{u}_j) \mathbf{P}_c \phi_{lmj} ds \right\}. \end{aligned}$$

We need to take a limit since $H_0 + e_j - e_l - e_m$ may not be invertible. Define

$$K_{lm}^j \equiv \lim_{r \rightarrow 0+} \frac{1}{H_0 - e_l - e_m + e_j - ri} \mathbf{P}_c. \quad (4.12)$$

$$\xi_{lm}^j \equiv -K_{lm}^j \mathbf{P}_c \phi_{lmj}. \quad (4.13)$$

K_{lm}^j are bounded operators in $\mathcal{B}(L_r^2, L_{-r'}^2)$ with $r, r' > 1/2$ and $r + r' > 2$, see [8]. Hence we have $\xi_{lm}^j \in L_{-r'}^2$. We may have $\xi_{lm}^j \notin L^2$ and $\text{Im } \xi_{lm}^j \neq 0$ only if

$$e_j - e_l - e_m < 0, \quad (4.14)$$

in particular $j < l, m$. We can now rewrite the above integral as

$$\begin{aligned} & x_l x_m \bar{x}_j(t) \xi_{lm}^j - x_l x_m \bar{x}_j(0) e^{-iH_0 t} \xi_{lm}^j \\ & - \int_0^t e^{-iH_0(t-s)} e^{i(-e_l - e_m + e_j)s} \frac{d}{ds} (u_l u_m \bar{u}_j) \xi_{lm}^j ds. \end{aligned}$$

Note that the choice $r \rightarrow 0+$, instead of $r \rightarrow 0-$, ensures the local decay of $e^{-iH_0 t} \xi_{lm}^j = -e^{-iH_0 t} K_{lm}^j \mathbf{P}_c \phi_{lmj}$ and of the last integral, see Lemma 2.3.

Summing these terms over l, m, j , we can decompose $\xi(t)$ as

$$\xi(t) = \xi^{(2)}(t) + \xi^{(3)}(t) = \xi^{(2)} + \left(\xi_1^{(3)} + \dots + \xi_5^{(3)} \right), \quad (4.15)$$

where $\xi^{(2)}(t)$ is the main part of $\xi(t)$,

$$\xi^{(2)}(t) = \sum_{l,m,j=0}^N x_l x_m \bar{x}_j(t) \xi_{lm}^j, \quad (4.16)$$

and $\xi^{(3)}(t) = \xi_1^{(3)} + \dots + \xi_5^{(3)}$ is the rest,

$$\begin{aligned} \xi_1^{(3)}(t) &= e^{-iH_0 t} \xi_0, \\ \xi_2^{(3)}(t) &= -e^{-iH_0 t} \xi^{(2)}(0), \\ \xi_3^{(3)}(t) &= - \int_0^t e^{-iH_0(t-s)} \mathbf{P}_c \sum_{l,m,j=0}^N e^{i(-e_l - e_m + e_j)s} \frac{d}{ds} (u_l u_m \bar{u}_j) \xi_{lm}^j ds, \\ \xi_4^{(3)}(t) &= \int_0^t e^{-iH_0(t-s)} \mathbf{P}_c i^{-1} (G - G_3 - \lambda \xi^2 \bar{\xi}) ds, \\ \xi_5^{(3)}(t) &= \int_0^t e^{-iH_0(t-s)} \mathbf{P}_c i^{-1} (\lambda \xi^2 \bar{\xi}) ds. \end{aligned} \quad (4.17)$$

We single out $\lambda |\xi|^2 \xi$ since it is the only non-local term in $G - G_3$.

We have the following estimates for nonlinear terms.

Lemma 4.1 *Suppose, for a fixed time t , for some $n \leq n_0$,*

$$|x_k(t)| \leq 2n, \quad \|\xi(t)\|_{L_{\text{loc}}^2 \cap L^5} \leq 2n, \quad \|\xi(t)\|_{L^2} \ll 1. \quad (4.18)$$

We have

$$\begin{aligned} \|G_7\|_{L_{\text{loc}}^1} &\lesssim n \|\xi\|_{L_{\text{loc}}^2}^2 + \|\xi\|_{L_{\text{loc}}^2}^{4/3} \|\xi\|_{L^5}^{5/3}, \\ \|G_5 + G_7\|_{L_{\text{loc}}^1} &\lesssim n^2 \|\xi\|_{L_{\text{loc}}^2}, \\ \|G\|_{L_{\text{loc}}^1} + \max_k |\dot{u}_k| &\lesssim n^3, \\ \|G_5 + G_7 - \lambda \xi^2 \bar{\xi}\|_{L^1 \cap L^{5/4}} &\lesssim n^2 \|\xi\|_{L_{\text{loc}}^2}. \end{aligned} \quad (4.19)$$

PROOF: For the nonlocal term $\lambda \xi^2 \bar{\xi}$ we have, using Hölder inequality,

$$\|\lambda \xi^2 \bar{\xi}\|_{L^{5/4}} \lesssim \|\xi\|_{L^2}^{2/3} \|\xi\|_{L^5}^{7/3}, \quad \|\lambda \xi^2 \bar{\xi}\|_{L^1} \lesssim \|\xi\|_{L^2}^{4/3} \|\xi\|_{L^5}^{5/3}. \quad (4.20)$$

Similarly, $\|\lambda \xi^2 \bar{\xi}\|_{L^1_{\text{loc}}} \lesssim \|\xi\|_{L^2_{\text{loc}}}^{4/3} \|\xi\|_{L^5}^{5/3}$. Using the definition (4.9) of G_7 , we have

$$\|G_7\|_{L^1_{\text{loc}}} \lesssim n \|\xi\|_{L^2_{\text{loc}}}^2 + \|\xi\|_{L^2_{\text{loc}}}^{4/3} \|\xi\|_{L^5}^{5/3}.$$

By (4.7) and (4.8), we have $\|G_3\| \lesssim n^3$ and $\|G_5\| \lesssim n^2 \|\xi\|_{L^2_{\text{loc}}}$. By (4.18),

$$\|G_5 + G_7\|_{L^1_{\text{loc}}} \lesssim n^2 \|\xi\|_{L^2_{\text{loc}}} + n \|\xi\|_{L^2_{\text{loc}}}^2 + \|\xi\|_{L^2_{\text{loc}}}^{4/3} \|\xi\|_{L^5}^{5/3} \lesssim n^2 \|\xi\|_{L^2_{\text{loc}}},$$

and $\|G\|_{L^1_{\text{loc}}} \leq \|G_3\|_{L^1_{\text{loc}}} + \|G_5 + G_7\|_{L^1_{\text{loc}}} \lesssim n^3 + n^2 \|\xi\|_{L^2_{\text{loc}}} \lesssim n^3$. Since $|\dot{u}_k| \leq \|G\|_{L^1_{\text{loc}}}$ by (4.4), the estimate for $\dot{u}_k$ follows. Finally, $G_7 - \lambda \xi^2 \bar{\xi}$ is of the form $n \xi^2$ and

$$\|G_7 - \lambda \xi^2 \bar{\xi}\|_{L^1 \cap L^{5/4}} \lesssim n \|\xi\|_{L^2_{\text{loc}} \cap L^5}^{2/3} \|\xi\|_{L^2_{\text{loc}}}^{4/3} \lesssim n^2 \|\xi\|_{L^2_{\text{loc}}}$$

by (4.18). Since $\|G_5\|_{L^1 \cap L^{5/4}} \lesssim n^2 \|\xi\|_{L^2_{\text{loc}}}$, the last estimate follows. **Q.E.D.**

4.3 Normal forms for equations of bound states

Recall we write $x_k(t) = e^{-ie_k t} u_k(t)$. In this subsection we derive the normal form for the equations of $\dot{u}_k$, where terms of different phase factors are removed.

Lemma 4.2 (Normal form) *Suppose for some $n \leq n_0$,*

$$|x_k(t)| \leq 2n, \quad \|\xi(t)\|_{L^2_{\text{loc}} \cap L^5} \leq 2n, \quad \|\xi(t)\|_{L^2} \ll 1. \quad (4.21)$$

There are perturbations $\mu_k(t)$ of $u_k(t)$, $k = 0, 1, \dots, N$, to be defined in (4.39), satisfying

$$|u_k(t) - \mu_k(t)| \leq Cn^3, \quad (4.22)$$

such that

$$\dot{\mu}_k = \sum_{l=0}^N c_l^k |\mu_l|^2 \mu_k + \sum_{a,b=0}^N d_{ab}^k |\mu_a|^2 |\mu_b|^2 \mu_k + g_k. \quad (4.23)$$

Here g_k are error terms, to be defined in (4.40), satisfying

$$|g_k(t)| \lesssim n^7 + n^2 \|\xi^{(3)}\|_{L^2_{\text{loc}}} + n \|\xi\|_{L^2_{\text{loc}}}^2 + \|\xi\|_{L^2_{\text{loc}}}^{4/3} \|\xi\|_{L^5}^{5/3}. \quad (4.24)$$

All coefficients c_l^k and d_{ab}^k are bounded by one. The coefficients c_l^k are purely imaginary. The real parts of d_{ab}^k are given by

$$\text{Re } d_{ab}^k = (2 - \delta_a^b) \gamma_{ab}^k - 2(2 - \delta_k^b) \gamma_{kb}^a, \quad (4.25)$$

where, with $\phi_{abk} = \lambda \phi_a \phi_b \phi_k$,

$$\gamma_{ab}^k \equiv \text{Im} \left(\phi_{abk}, \frac{1}{H_0 - e_a - e_b + e_k - 0i} \mathbf{P}_c \phi_{abk} \right) \geq 0. \quad (4.26)$$

Note γ_{ab}^k may be positive only if $e_a - e_b + e_k < 0$, in particular $k < a, b$. By assumption A2 we have $\gamma_{kk}^0 \geq \gamma_0 > 0$ for $k > 0$. The first part of $\text{Re } d_{ab}^k$ is positive and the second part negative. If $k = 0$, $\gamma_{kb}^a = 0$ for all a, b , hence $\text{Re } d_{ab}^k \geq 0$. If $k = N$, $\gamma_{ab}^k = 0$ for all a, b , hence $\text{Re } d_{ab}^k \leq 0$. For intermediate k the sign of $\text{Re } d_{ab}^k$ depends on the relative size of these coefficients. Let $f_k = |\mu_k|^2$, $k = 0, \dots, N$. Since $\frac{d}{dt}|\mu|^2 = 2 \text{Re } \bar{\mu} \dot{\mu}$ and c_l^k are purely imaginary, we have

$$\dot{f}_k = \sum_{a,b=0}^N 2(\text{Re } d_{ab}^k) f_a f_b f_k + 2 \text{Re } \bar{\mu}_k g_k. \quad (4.27)$$

The example 1 in §1 follows by letting $N = 2$, $x = \mu_0$, $y = \mu_1$ and $z = \mu_2$.

PROOF: Recall (4.4) that

$$\dot{u}_k = i^{-1} e^{ie_k t} (\phi_k, G), \quad (k = 0, \dots, N). \quad (4.28)$$

We now proceed to derive the normal form in two steps.

Step 1 Integration of terms of order n^3 .

Substituting $G = G_3 + G_5 + G_7$ and the expression (4.10) for G_3 , we get

$$\begin{aligned} \dot{u}_k(t) &= \sum_{l,m,j=0}^N i^{-1} e^{ie_k t} x_l x_m \bar{x}_j(t) (\phi_k, \phi_{lmj}) + R_k \\ &= \sum_{l,m,j=0}^N i^{-1} e^{i(e_k - e_l - e_m + e_j)t} u_l u_m \bar{u}_j(t) (\phi_k, \phi_{lmj}) + R_k, \\ R_k &= i^{-1} e^{ie_k t} (\phi_k, G_5 + G_7)(t). \end{aligned} \quad (4.29)$$

By assumption A3,

$$e_k - e_l - e_m + e_j \neq 0, \quad (4.30)$$

unless the two sets $\{k, n\}$ and $\{l, m\}$ are the same. Hence

$$\dot{u}_k(t) = \sum_{l=0}^N c_l^k |u_l|^2 u_k + \sum_{(4.30)} i^{-1} e^{i(e_k - e_l - e_m + e_j)t} u_l u_m \bar{u}_j(t) (\phi_k, \phi_{lmj}) + R_k, \quad (4.31)$$

where c_l^k are purely imaginary constants defined by

$$c_l^k = 2i^{-1}(\phi_k, \phi_{ulk}), \quad (k \neq l); \quad c_k^k = i^{-1}(\phi_k, \phi_{kkk}).$$

Terms in the second group of (4.31) are oscillatory and can be integrated. Define

$$u_{k,1} = u_k - u_{k,1}^-,$$

where

$$u_{k,1}^-(t) = \sum_{(4.30)} \frac{i^{-1} e^{i(e_k - e_l - e_m + e_j)t}}{i(e_k - e_l - e_m + e_j)} u_l u_m \bar{u}_j(t) (\phi_k, \phi_{lmj}),$$

$$g_{k,1}(t) = - \sum_{(4.30)} \frac{i^{-1} e^{i(e_k - e_l - e_m + e_j)t}}{i(e_k - e_l - e_m + e_j)} \frac{d}{dt} (u_l u_m \bar{u}_j)(t) (\phi_k, \phi_{lmj}).$$

We have

$$\dot{u}_{k,1} = \dot{u}_k - \dot{u}_{k,1}^- = \sum_{l=0}^N c_l^k |u_l|^2 u_k + R_k + g_{k,1}.$$

We further define

$$g_{k,2} = \sum_{l=0}^N c_l^k \{ |u_l|^2 u_k - |u_{l,1}|^2 u_{k,1} \}.$$

We have

$$\dot{u}_{k,1} = \sum_{l=0}^N c_l^k |u_{l,1}|^2 u_{k,1} + i^{-1} e^{ie_k t} (\phi_k, G_5 + G_7) + g_{k,1} + g_{k,2}. \quad (4.32)$$

Step 2 Integration of terms of order n^5 .

We next integrate $O(n^5)$ terms. We first decompose $g_{k,1}$ and $g_{k,2}$ since they contain $O(n^5)$ terms. Using their explicit form we can decompose them as

$$g_{k,1} = e^{ie_k t} g_{k,1,5} + g_{k,1,7}, \quad (4.33)$$

$$g_{k,2} = e^{ie_k t} g_{k,2,5} + g_{k,2,7}, \quad (4.34)$$

where $g_{k,1,7}$ and $g_{k,2,7}$ are higher order terms with

$$|g_{k,1,7}| \lesssim n^2 \max_j |R_j|, \quad |g_{k,2,7}| \lesssim n^7. \quad (4.35)$$

$g_{k,1,5}$ and $g_{k,2,5}$ are explicit homogeneous polynomials of degree 5 of the form

$$C x_{l_1} x_{l_2} x_{l_3} \bar{x}_{l_4} \bar{x}_{l_5}, \quad (4.36)$$

where $l_j \in \{0, \dots, N\}$ and C is a *purely imaginary* coefficient. The claim on $g_{k,1,5}$ can be seen by substituting (4.29) into $\frac{d}{dt}(u_l u_m \bar{u}_j)$ in the definition of $g_{k,1}$. The claim on $g_{k,2,5}$ can be seen from the definitions of $g_{k,2}$ and $u_{k,1}^-$.

We next decompose $i^{-1} e^{ie_k t} (\phi_k, G_5)$. Recall $G_5 = \lambda \chi^2 \bar{\xi} + 2\lambda |\chi|^2 \xi$ with $\chi = \sum_{l=0}^N x_l \phi_l$. Using the decomposition $\xi = \xi^{(2)} + \xi^{(3)}$ with $\xi^{(2)} = \sum_{l,m,j=0}^N x_l x_m \bar{x}_j \xi_{lm}^j$, we can decompose

$$G_5 = G_{5,1} + G_{5,2} + G_{5,3},$$

where

$$\begin{aligned} G_{5,1} &= \lambda \chi^2 \sum_{l,m,j=0}^N \bar{x}_l \bar{x}_m x_j (-i \operatorname{Im} \xi_{lm}^j) + 2\lambda |\chi|^2 \sum_{l,m,j=0}^N x_l x_m \bar{x}_j (i \operatorname{Im} \xi_{lm}^j), \\ G_{5,2} &= \lambda \chi^2 \sum_{l,m,j=0}^N \bar{x}_l \bar{x}_m x_j (\operatorname{Re} \xi_{lm}^j) + 2\lambda |\chi|^2 \sum_{l,m,j=0}^N x_l x_m \bar{x}_j (\operatorname{Re} \xi_{lm}^j), \\ G_{5,3} &= \lambda \chi^2 \bar{\xi}^{(3)} + 2\lambda |\chi|^2 \xi^{(3)}. \end{aligned}$$

Recall that $\text{Im } \xi_{lm}^j \neq 0$ only if $e_j - e_l - e_m < 0$. The term $G_{5,3}$ will be shown to be smaller than $G_{5,1}$ and $G_{5,2}$. Both $G_{5,1}$ and $G_{5,2}$ are homogeneous polynomials of degree 5 of the form (4.36). However, the coefficients C of $G_{5,2}$ are real-valued L_{loc}^2 functions, while those of $G_{5,1}$ are purely imaginary.

We can now rewrite (4.32) as

$$\dot{u}_{k,1} = \sum_{l=0}^N c_l^k |u_{l,1}|^2 u_{k,1} + \tilde{R}_k + (i^{-1} e^{ie_k t} (\phi_k, G_{5,3} + G_7) + g_{k,1,7} + g_{k,2,7}),$$

where $\tilde{R}_k$ consist of terms of order $O(n^5)$,

$$\tilde{R}_k = e^{ie_k t} \{ i^{-1} (\phi_k, G_{5,1} + G_{5,2}) + g_{k,1,5} + g_{k,2,5} \}. \quad (4.37)$$

Inside the bracket are polynomials in x_l and $\bar{x}_l$, $l = 0, \dots, N$. The only terms with real coefficients are those from $G_{5,1}$. As in Step 1, the above expression can be separated into two groups: those with zero phase factor and those with non-zero phase factors. The phase factor of a typical term in the above expression is of the form

$$e_k - e_{l_1} - e_{l_2} - e_{l_3} + e_{l_4} + e_{l_5},$$

where $l_j \in \{0, \dots, N\}$. To get zero, we must have the form $e_k - e_k - e_a - e_b + e_a + e_b$ for some a, b by assumption A3. Thus a typical term in the first group is of the form $C e^{ie_k t} x_k x_a x_b \bar{x}_a \bar{x}_b = C |u_a|^2 |u_b|^2 u_k$.

We can integrate by parts those terms with non-zero phase factors as in Step 1. The other terms remain. Hence we can rewrite $\tilde{R}_k$ as

$$\tilde{R}_k = \sum_{a,b=0}^N d_{ab}^k |u_a|^2 |u_b|^2 u_k + \frac{d}{dt} (u_{k,2}^-) + g_{k,3},$$

for some order one constants d_{ab}^k , some explicit homogeneous polynomials $u_{k,2}^-$ of degree 5, and remainder terms $g_{k,3}$ with

$$|g_{k,3}| \lesssim n^4 \max_j |\dot{u}_j|. \quad (4.38)$$

We will identify $\text{Re } d_{ab}^k$ in a moment. Define

$$\mu_k \equiv u_{k,1} - u_{k,2}^- = u_k - u_{k,1}^- - u_{k,2}^-. \quad (4.39)$$

We have

$$\begin{aligned} \dot{\mu}_k &= \dot{u}_{k,1} - \dot{u}_{k,2}^- \\ &= \sum_{l=0}^N c_l^k |u_{l,1}|^2 u_{k,1} + \sum_{a,b=0}^N d_{ab}^k |u_a|^2 |u_b|^2 u_k + g_{k,3} \\ &\quad + (i^{-1} e^{ie_k t} (\phi_k, G_{5,3} + G_7) + g_{k,1,7} + g_{k,2,7}) \\ &= \sum_{l=0}^N c_l^k |\mu_l|^2 \mu_k + \sum_{a,b=0}^N d_{ab}^k |\mu_a|^2 |\mu_b|^2 \mu_k + g_k, \end{aligned}$$

where

$$g_k = i^{-1} e^{ie_k t} (\phi_k, G_{5,3} + G_7) + g_{k,1,7} + g_{k,2,7} + g_{k,3} + g_{k,4}, \quad (4.40)$$

$$g_{k,4} = \sum_{l=0}^N c_l^k \{ |u_{l,1}|^2 u_{k,1} - |\mu_l|^2 \mu_k \} + \sum_{a,b=0}^N d_{ab}^k \{ |u_a|^2 |u_b|^2 u_k - |\mu_a|^2 |\mu_b|^2 \mu_k \}.$$

We have $|g_{k,5}| \lesssim n^7$. Collecting estimates, using the explicit form of $G_{5,3}$, $|R_k| \lesssim \|G_5 + G_7\|_{L_{\text{loc}}^1}$, and Lemma 4.1, we get

$$\begin{aligned} |g_k| &\lesssim \|G_{5,3}\|_{L_{\text{loc}}^1} + \|G_7\|_{L_{\text{loc}}^1} + n^2 \max_j |R_j| + n^7 + n^4 \max_j |\dot{u}_j| + n^7 \\ &\lesssim n^7 + n^2 \|\xi^{(3)}\|_{L_{\text{loc}}^2} + n^4 \|\xi\|_{L_{\text{loc}}^2} + n \|\xi\|_{L_{\text{loc}}^2}^2 + \|\xi^3\|_{L_{\text{loc}}^1}. \end{aligned}$$

Using $\|\xi\|_{L_{\text{loc}}^2} \lesssim n^3 + \|\xi^{(3)}\|_{L_{\text{loc}}^2}$, we can remove $n^4 \|\xi\|_{L_{\text{loc}}^2}$ and get (4.24).

We have noted that c_l^k are purely imaginary. We now compute the real parts of d_{ab}^k . From the previous discussions on $g_{k,1,5}$, $g_{k,2,5}$, $G_{5,1}$ and $G_{5,2}$, we know that the only contribution to $\text{Re } d_{ab}^k$ is from $i^{-1} e^{ie_k t} (\phi_k, G_{5,1})$. We have

$$\begin{aligned} i^{-1} e^{ie_k t} (\phi_k, G_{5,1}) &= \sum_{a,b,l,m,j} -e^{ie_k t} x_a x_b \bar{x}_l \bar{x}_m x_j (\phi_k, \lambda \phi_a \phi_b \text{Im } \xi_{lm}^j) \\ &\quad + \sum_{a,b,l,m,j} 2e^{ie_k t} x_a \bar{x}_b x_l x_m \bar{x}_j (\phi_k, \lambda \phi_a \phi_b \text{Im } \xi_{lm}^j), \end{aligned}$$

and we are interested in those terms of the form $C e^{ie_k t} |u_a|^2 |u_b|^2 u_k$. As mentioned, $\text{Im } \xi_{lm}^j \neq 0$ only if $e_j - e_l - e_m < 0$, in particular $j \neq l, m$ (not sufficient though). Therefore, to get products in the first summation we need $j = k$ and the two sets $\{l, m\}$ and $\{a, b\}$ are the same; in the second summation we need $j = a$ and the two sets $\{l, m\}$ and $\{k, b\}$ are the same. Hence $\sum_{a,b=0}^N (\text{Re } d_{ab}^k) |u_a|^2 |u_b|^2 u_k$ is equal to

$$\begin{aligned} &\sum_{a,b=0}^N -(2 - \delta_a^b) |x_a|^2 |x_b|^2 u_k (\phi_k, \lambda \phi_a \phi_b \text{Im } \xi_{ab}^k) \\ &+ \sum_{a,b=0}^N 2(2 - \delta_k^b) |x_a|^2 |x_b|^2 u_k (\phi_k, \lambda \phi_a \phi_b \text{Im } \xi_{kb}^a). \end{aligned}$$

Note we have the factors $(2 - \delta_a^b)$ and $(2 - \delta_k^b)$ since we have two choices for assigning l and m if $a \neq b$ (or $k \neq b$). Recall $\lambda \phi_a \phi_b \phi_k = \phi_{abk}$, $\xi_{lm}^j = -K_{lm}^j \mathbf{P}_c \phi_{lmj}$, and $\gamma_{lm}^j = (\phi_{lmj}, \text{Im } K_{lm}^j \mathbf{P}_c \phi_{lmj})$. We conclude

$$\begin{aligned} \text{Re } d_{ab}^k &= -(2 - \delta_a^b) (\phi_{abk}, \text{Im } \xi_{ab}^k) + 2(2 - \delta_k^b) (\phi_{kba}, \text{Im } \xi_{kb}^a) \\ &= (2 - \delta_a^b) \gamma_{ab}^k - 2(2 - \delta_k^b) \gamma_{kb}^a. \end{aligned}$$

Q.E.D.

4.4 Main estimates

In this subsection we prove estimates in the transition regime for the solution $\psi(t)$. The following proposition is the main result of this section. It proves the same result of Theorem 1.1 under weaker assumptions (4.41)–(4.42) below. Note that the assumption $\|\xi_0\|_Y \leq n/2$ in (1.9) of Theorem 1.1 implies, using Lemma 2.3,

$$\|e^{-itH_0}\xi_0\|_{L^5} \lesssim n(1+t)^{-9/10}, \quad \|e^{-itH_0}\xi_0\|_{L^2_{\text{loc}}} \lesssim n(1+t)^{-3/2}.$$

Hence it implies (4.42). Eq. (4.42) is motivated by (4.46) and [22], and may be more convenient for future application when we study the asymptotic profiles of all small solutions. See [22] for two bound states case. Also note that, by (4.41) below, we have $\|\psi(t)\|_{L^2} = \|\psi_0\|_{L^2} \ll 1$ and hence $\max_k |x_k(t)|, \|\xi(t)\|_{L^2} \ll 1$. However, we do not assume any bound of $\|\psi_0\|_{L^2}$ in terms of n . This is essential for future application.

Notice that we define t_2 by (4.45) to be the time when the size of the excited states decays to the order $n^{1+\sigma}$. This is more than sufficient since we only need it to be smaller than $\varepsilon_0 n$ in order to apply Theorem 3.1.

Proposition 4.3 *Assume the assumptions A0–A3 given in §1. There is a small constant $n_0 > 0$ such that the following hold. Suppose $\psi(t) = \psi(t, x)$ is a solution of (1.1) with the initial data $\psi(0) = \psi_0 = x_0^0 \phi_0 + \cdots + x_N^0 \phi_N + \xi_0$ satisfying, for some $n \leq n_0$, $0 < \sigma < 1/10$,*

$$\begin{aligned} \frac{3}{4}n^2 &\leq |x_0^0|^2 + \cdots + |x_N^0|^2 \leq n^2, & \|\psi_0\|_{H^1} &\ll 1, \\ |x_0^0| &\geq n^{3-\epsilon}, & \left(\sum_{k=1}^N |x_k^0|^2\right)^{1/2} &\geq 2|x_0^0|^{1+\sigma}, \end{aligned} \tag{4.41}$$

and for some $t_* \in [1, n^{-4-2\sigma}]$, for all $t \geq 0$,

$$\begin{aligned} \|e^{-itH_0}\xi_0\|_{L^5} &\leq \Lambda_5(t) \equiv n(1+t)^{-9/10} + Cn^3 t_* (t_* + t)^{-9/10}, \\ \|e^{-itH_0}\xi_0\|_{L^2_{\text{loc}}} &\leq \Lambda(t) \equiv n(1+t)^{-3/2} + Cn^3 \frac{t_*}{t_* + t} (1+t)^{-1/2}. \end{aligned} \tag{4.42}$$

Let $t_0 = n^{-2}$. There exist $t_2 \geq t_1 \geq t_0$ such that

$$t_1 \leq C_4 n^{-4} \log \frac{1}{n}, \quad C_4^{-1} n^{-4} \leq t_2 \leq C_4 n^{-4-2\sigma}, \tag{4.43}$$

for some constant $C_4 > 1$ independent of n , and we have, for $0 \leq t \leq t_2$,

$$\begin{aligned} |x_0(t)| &\geq \frac{3}{4} \sup_{0 \leq s \leq t} |x_0(s)|, \\ |x_k(t)| &\leq \frac{5}{4}n, \quad (k = 0, \dots, N), \\ \|\xi(t)\|_{L^5} &\leq C_5 n^3 t^{1/10} + \frac{5}{4} \Lambda_5(t), \\ \|\xi(t)\|_{L^2_{\text{loc}}} &\leq C_5 n^3 + \frac{5}{4} \Lambda(t), \\ \|\xi^{(3)}(t)\|_{L^2_{\text{loc}}} &\leq C_5 n^5, \quad \text{for } t \geq t_0, \end{aligned} \tag{4.44}$$

where C_5 is an explicit constant. We also have, at $t = t_1$ and $t = t_2$,

$$|x_0(t_1)|^2, |x_0(t_2)|^2 \geq \frac{3}{4} 2^{-2-N} n^2, \quad \left(\sum_{k=1}^N |x_k(t_2)|^2 \right)^{1/2} \leq |x_0(t_2)|^{1+\sigma}. \quad (4.45)$$

Moreover, we have the following out-going estimates for $\xi(t_2)$: For $\tau \geq 0$,

$$\begin{aligned} \|e^{-i\tau H_0} \xi(t_2)\|_{L^5} &\leq C n^3 (t_* + t_2) (t_2 + \tau)^{-9/10}, \\ \|e^{-i\tau H_0} \xi(t_2)\|_{L^2_{\text{loc}}} &\leq C n^3 \frac{t_* + t_2}{t_2 + \tau} (1 + \tau)^{-1/2}. \end{aligned} \quad (4.46)$$

PROOF: We will prove estimates (4.43)–(4.45) using a continuity argument. Hence we can assume the following weaker estimates: For $0 \leq t \leq t_2$:

$$\begin{aligned} |x_0(t)| &\geq \frac{1}{2} \sup_{0 \leq s \leq t} |x_0(s)|, \\ |x_k(t)| &\leq \frac{3}{2} n, \quad (k = 0, \dots, N), \\ \|\xi(t)\|_{L^5} &\leq 2C_5 n^3 t^{1/10} + 2\Lambda_5(t), \\ \|\xi(t)\|_{L^2_{\text{loc}}} &\leq 2C_5 n^3 + 2\Lambda(t), \\ \|\xi^{(3)}(t)\|_{L^2_{\text{loc}}} &\leq 2C_5 n^5, \quad \text{for } t \geq t_0. \end{aligned} \quad (4.47)$$

If we can prove (4.43)–(4.45) assuming (4.47), then (4.43)–(4.45) hold true for all $t \in [0, t_2]$ since (4.47) is never violated. By definition we have $\Lambda_5(t), \Lambda(t) \leq n$. By (4.47) we have $\|\xi(t)\|_{L^2_{\text{loc}} \cap L^5} \leq 2n$. Hence the assumptions of Lemmas 4.1 and Lemma 4.2 are satisfied. Since $\|\psi(t)\|_{L^2} = \|\psi_0\|_{L^2} \ll 1$, we have $\|\xi(t)\|_{L^2} \ll 1$. Also, since $t \leq t_2 \leq C_4 n^{-4-2\sigma}$,

$$\|\xi\|_{L^5} \lesssim n^3 t_2^{1/10} + n(1+t)^{-9/10} + C n^3 t_*^{1/10} \lesssim n^{13/5-\sigma/5} + n(1+t)^{-9/10}, \quad (4.48)$$

$$\|\xi\|_{L^2_{\text{loc}}} \lesssim n^3 + n(1+t)^{-3/2} + C n^3 (1+t)^{-1/2} \lesssim n^3 + n(1+t)^{-3/2}. \quad (4.49)$$

We now estimate $\xi(t)$. Recall (4.5),

$$\xi(t) = e^{-itH_0} \xi_0 + \int_0^t e^{-i(t-s)H_0} \mathbf{P}_c i^{-1} G(s) ds.$$

Note $\|G\|_{L^{5/4}} \lesssim \|G_3\|_{L^{5/4}} + \|G - G_3 - \lambda \xi^2 \bar{\xi}\|_{L^{5/4}} + \|\lambda \xi^2 \bar{\xi}\|_{L^{5/4}}$ and $\|G_3\|_{L^{5/4}} \leq C n^3$. Using Lemma 4.1, (4.48) and (4.49),

$$\|G - G_3 - \lambda \xi^2 \bar{\xi}\|_{L^1 \cap L^{5/4}} \lesssim n^2 \|\xi\|_{L^2_{\text{loc}}} \lesssim n^2 (n^3 + n(1+s)^{-3/2}), \quad (4.50)$$

$$\|\lambda \xi^2 \bar{\xi}\|_{L^{5/4}} \lesssim \|\xi\|_{L^2}^{2/3} \|\xi\|_{L^5}^{7/3} \leq o(1) [n^{13/5-\sigma/5} + n(1+s)^{-9/10}]^{7/3}. \quad (4.51)$$

Hence $\|G(s)\|_{L^{5/4}} \lesssim n^3 + n^{7/3} \langle s \rangle^{-21/10}$ and

$$\begin{aligned} \|\xi(t)\|_{L^5} &\leq \|e^{-itH_0}\xi_0\|_{L^5} + C \int_0^t |t-s|^{-9/10} \|G(s)\|_{L^{5/4}} ds \\ &\leq \Lambda_5(t) + C \int_0^t |t-s|^{-9/10} [n^3 + n^{7/3} \langle s \rangle^{-21/10}] ds \leq \frac{5}{4}\Lambda_5(t) + Cn^3 t^{1/10}. \end{aligned}$$

Hence $\|\xi(t)\|_{L^5}$ is estimated. For the L_{loc}^2 norm, we use the decomposition (4.15) that $\xi(t) = \xi^{(2)} + \xi^{(3)}$ with $\xi^{(3)} = \sum_{j=1}^5 \xi_j^{(3)}$. We have $\|\xi^{(2)}\|_{L_{\text{loc}}^2} \leq Cn^3$ by its explicit form. By (4.42) for ξ_0 we have

$$\left\| \xi_1^{(3)}(t) \right\|_{L_{\text{loc}}^2} \leq \Lambda(t).$$

By the singular decay estimate (2.5) in Lemma 2.3 and the definition (4.13) of ξ_{lm}^j , we have

$$\left\| \xi_2^{(3)}(t) \right\|_{L_{\text{loc}}^2} \leq Cn^3(1+t)^{-3/2} \leq Cn^2\Lambda(t).$$

Using (2.5), (4.13) again and the bound of $\max_j |\dot{u}_j|$ in Lemma 4.1, we have

$$\left\| \xi_3^{(3)}(t) \right\|_{L_{\text{loc}}^2} \leq C \int_0^t \langle t-s \rangle^{-3/2} n^2 \max_j |\dot{u}_j| ds \leq C \int_0^t \langle t-s \rangle^{-3/2} n^5 ds \leq Cn^5.$$

For $\xi_4^{(3)}(t)$, bounding its integrand by either L^∞ or L^5 -norm and using (4.50), we have

$$\begin{aligned} \left\| \xi_4^{(3)}(t) \right\|_{L_{\text{loc}}^2} &\lesssim \int_0^t \min \{ |t-s|^{-3/2}, |t-s|^{-9/10} \} \|(G - G_3 - \lambda \xi^2 \bar{\xi})(s)\|_{L^1 \cap L^{5/4}} ds \\ &\lesssim \int_0^t \min \{ |t-s|^{-3/2}, |t-s|^{-9/10} \} n^2 (n^3 + n(1+s)^{-3/2}) ds \\ &\lesssim n^5 + n^2 \Lambda(t). \end{aligned}$$

Finally we give two estimates for $\xi_5^{(3)}$. By Hölder inequality and (4.48),

$$\begin{aligned} \|\xi^3\|_{L^1 \cap L^{5/4}} &\leq \|\xi\|_{L^2 \cap L^5}^{4/3} \|\xi\|_{L^5}^{5/3} \leq o(1)(n^{13/5-\sigma/5} + n(1+s)^{-9/10})^{5/3} \\ &\lesssim n^{13/3-\sigma/3} + n^{5/3}(1+s)^{-3/2}. \end{aligned} \tag{4.52}$$

The same estimate for $\xi_4^{(3)}$ shows

$$\left\| \xi_5^{(3)}(t) \right\|_{L_{\text{loc}}^2} \lesssim n^{13/3-\sigma/3} + n^{2/3}\Lambda(t).$$

Summing the estimates we conclude

$$\|\xi\|_{L_{\text{loc}}^2} \leq \|\xi^{(2)}\|_{L_{\text{loc}}^2} + \sum_{j=1}^5 \left\| \xi_j^{(3)} \right\|_{L_{\text{loc}}^2} \leq C_5 n^3 + \frac{5}{4}\Lambda(t).$$

We now estimate $\xi_5^{(3)}$ in another way. We have

$$\begin{aligned} \|\xi^2 \bar{\xi}\|_{L^{6/5} \cap L^{5/4}} &\leq \|\xi\|_{L^2 \cap L^5}^{7/9} \|\xi\|_{L^5}^{20/9} \leq o(1)(n^{13/5-\sigma/5} + n(1+s)^{-9/10})^{20/9} \\ &\leq n^{52/9-4\sigma/9} + n^{20/9}(1+s)^{-2}. \end{aligned}$$

Bounding the integrand of $\xi_5^{(3)}(t)$ by either L^6 or L^5 norm we have

$$\begin{aligned} \|\xi_5^{(3)}(t)\|_{L_{\text{loc}}^2} &\lesssim \int_0^t \min\{|t-s|^{-1}, |t-s|^{-9/10}\} \|\lambda \xi^2 \bar{\xi}(s)\|_{L^{6/5} \cap L^{5/4}} ds \\ &\lesssim \int_0^t \min\{|t-s|^{-1}, |t-s|^{-9/10}\} (n^{52/9-4\sigma/9} + n^{20/9}(1+s)^{-2}) ds \\ &\lesssim n^{52/9-4\sigma/9} \log(1+t_2) + n^{20/9}(1+t)^{-1} \log(1+t_2). \end{aligned}$$

Summing the estimates and using $t_2 \leq C_4 t^{-4-2\sigma}$, we get

$$\|\xi^{(3)}\|_{L_{\text{loc}}^2} \leq \sum_{j=1}^5 \|\xi_j^{(3)}\|_{L_{\text{loc}}^2} \leq Cn^5 + \frac{5}{4}\Lambda(t) + n^{20/9}(1+t)^{-1} \log(1+t_2).$$

Clearly we have $\|\xi^{(3)}(t)\|_{L_{\text{loc}}^2} \leq Cn^5$ for $t > t_0 = n^{-2}$.

We now estimate the error term $g_k(t)$ in the equation (4.23) of u_k . Recall (4.24),

$$|g_k(t)| \lesssim n^7 + n^2 \|\xi^{(3)}\|_{L_{\text{loc}}^2} + n \|\xi\|_{L_{\text{loc}}^2}^2 + \|\xi\|_{L_{\text{loc}}^2}^{4/3} \|\xi\|_{L^5}^{5/3}.$$

Using (4.47)–(4.49), $\Lambda_5(t), \Lambda(t) \leq n$, and the computation in (4.52),

$$\begin{aligned} |g_k(t)| &\lesssim n^7 + n^2 \|\xi^{(3)}\|_{L_{\text{loc}}^2} + n(n^3 + \Lambda)^2 + (n^3 + \Lambda)^{4/3} (n^{13/3-\sigma/3} + n^{2/3}\Lambda(s)) \\ &\lesssim n^7 + n^2 \|\xi^{(3)}\|_{L_{\text{loc}}^2} + n^{14/3-\sigma/3} \Lambda + n\Lambda^2. \end{aligned}$$

For $t \leq t_0 = n^{-2}$ we use the first estimate of $\|\xi_5^{(3)}\|_{L_{\text{loc}}^2}$ and we have $\|\xi^{(3)}\|_{L_{\text{loc}}^2} \leq \sum_{j=1}^5 \|\xi_j^{(3)}\|_{L_{\text{loc}}^2} \lesssim n^{13/3-\sigma/3} + \Lambda$. Hence

$$|g_k(t)| \lesssim n^{6+1/3-\sigma/3} + n^2 \Lambda, \quad (t \in [0, t_0]). \quad (4.53)$$

For $t \geq t_0$ we have $\|\xi^{(3)}(t)\|_{L_{\text{loc}}^2} \leq Cn^5$ and $\Lambda(t) \lesssim n^2$. Thus

$$|g_k(t)| \lesssim n^7, \quad (t \in [t_0, t_2]). \quad (4.54)$$

We now estimate bound states. As illustrated in Example 1 of §1, some bound states may grow at intermediate time. We will use the following monotonicity properties. Recall that $f_k = |\mu_k|^2$, $k = 0, \dots, N$, satisfy (4.27). Motivated by (1.19), we consider the following functions:

$$f(t) = f_0(t), \quad g(t) = \sum_{k=1}^N f_k, \quad g_+(t) = \sum_{k=1}^N 2^{-k} f_k. \quad (4.55)$$

By (4.41), (4.42) and (4.22) we have

$$f(0) \geq 2^{-1}n^{6-2\epsilon}, \quad 2n^2 \geq f(0) + g(0) \geq 2^{-1}n^2, \quad f(0) + g_+(0) \geq 2^{-1-N}n^2, \quad (4.56)$$

if n is sufficiently small. Although g and g_+ are comparable, $(f + g)(t)$ is almost monotone decreasing while $(f + g_+)(t)$ is almost monotone increasing in the following sense:

$$\begin{aligned} \frac{d}{dt}(f + g)(t) &\leq 2(N + 1) \max_k |\bar{\mu}_k g_k|, \\ \frac{d}{dt}(f + g_+)(t) &\geq -2(N + 1) \max_k |\bar{\mu}_k g_k|. \end{aligned} \quad (4.57)$$

We now prove (4.57). By (4.25) and (4.27),

$$\begin{aligned} \frac{d}{dt}(f + g) - \sum_{k=0}^N 2 \operatorname{Re} \bar{\mu}_k g_k &= \sum_{a,b,k=0}^N 2(\operatorname{Re} d_{ab}^k) f_a f_b f_k \\ &= \sum_{a,b,k=0}^N 2[(2 - \delta_a^b) \gamma_{ab}^k - 2(2 - \delta_k^b) \gamma_{kb}^a] f_a f_b f_k. \end{aligned} \quad (4.58)$$

Switching a and k in the terms with minus sign, we see that the above sum is non-positive. Similarly,

$$\begin{aligned} \frac{d}{dt}(f + g_+) - \sum_{k=0}^N 2^{1-k} \operatorname{Re} \bar{\mu}_k g_k &= \sum_{a,b,k=0}^N 2^{1-k} (\operatorname{Re} d_{ab}^k) f_a f_b f_k \\ &= \sum_{a,b,k=0}^N 2^{1-k} [(2 - \delta_a^b) \gamma_{ab}^k - 2(2 - \delta_k^b) \gamma_{kb}^a] f_a f_b f_k \\ &= \sum_{a,b,k=0}^N [2^{1-k} (2 - \delta_a^b) \gamma_{ab}^k - 2^{2-a} (2 - \delta_a^b) \gamma_{ab}^k] f_a f_b f_k. \end{aligned}$$

In the last line we have switched a and k for terms with minus sign. If $k \geq a$ then $\gamma_{ab}^k = 0$. Hence $2^{1-k} \geq 2^{2-a}$ for nonzero terms and the above sum is non-negative. We have shown (4.57). We now estimate the bound states in three steps.

Step 1. Initial layer.

In this period the dispersive part disperses away so much that it becomes negligible locally. The time it takes is less than $t_0 = n^{-2}$, which is not long enough to change the magnitudes of the bound states. Explicitly, by (4.27), (4.47) and (4.53) we have

$$|\dot{f}_k(s)| \leq Cn^4\alpha^2 + \alpha(n^{6+1/3-\sigma/3} + n^2\Lambda(s)),$$

for $k = 0, 1, \dots, N$, where $\alpha = \sqrt{f_k(0)}$. Thus, for $t \in [0, t_0]$,

$$\begin{aligned} |f_k(t) - f_k(0)| &\lesssim \int_0^t n^4\alpha^2 + \alpha(n^{6+1/3-\sigma/3} + n^2\Lambda(s)) ds \\ &\lesssim n^4\alpha^2 t_0 + \alpha n^{6+1/3-\sigma/3} t_0 + \alpha n^2(n + n^3\sqrt{t_0}) \\ &\lesssim n^2\alpha^2 + n^3\alpha \lesssim n^\epsilon\alpha^2 + n^{6-\epsilon}. \end{aligned}$$

Since $f_k(0) \geq n^{6-2\epsilon}$, we get $|f_k(t) - f_k(0)| \ll f_k(0)$.

Step 2. Transition regime (i).

After time $t = t_0$ most mass of the dispersive wave is far away and has no effect on the local dynamics. Two possible situations can occur: We either have $f(t_0) \geq 2^{-2-N}n^2$ or the opposite. In the first case we define $t_1 = t_0$ and jump to next step. We now focus on the second case. In the second case the ground state begins to grow exponentially until it is of order n . The time it takes is of order $n^{-4} \log n^{-1}$. Define

$$t_1 \equiv \inf_{t \geq t_0} \{t : f_0(t) \geq 2^{-2-N}n^2\}. \quad (4.59)$$

By assumption $t_1 > t_0$. We want to show that

$$t_0 \leq t_1 \leq t'_1 \equiv t_0 + 2^{8+2N}N\gamma_0^{-1} n^{-4} \log \frac{1}{n}. \quad (4.60)$$

Suppose (4.60) fails, that is, $f_0(t) < 2^{-2-N}n^2$ for all $t \leq t'_1$. We have by (4.56), (4.54) and (4.57), for all $t \in [t_0, t'_1]$,

$$\begin{aligned} g_+(t) &\geq (f_0 + g_+)(t_0) - f_0(t) - \int_{t_0}^t \max_k |\mu_k| |g_k| ds \\ &\geq 2^{-1-N}n^2 - 2^{-2-N}n^2 - Cnn^7 \geq 2^{-3-N}n^2. \end{aligned}$$

In particular, the coefficient of the linear term in (4.27) of $\dot{f}_0$ has a lower bound,

$$\begin{aligned} \sum_{a,b=0}^N 2(\operatorname{Re} d_{ab}^0) f_a f_b &= \sum_{a,b=0}^N 2(2 - \delta_a^b) \gamma_{ab}^0 f_a f_b \geq \sum_{a=1}^N 2\gamma_0 f_a^2 \geq 2\gamma_0 N^{-1} g^2 \\ &\geq 2\gamma_0 N^{-1} g_+^2 \geq 2^{-5-2N} \gamma_0 N^{-1} n^4. \end{aligned} \quad (4.61)$$

Since $\sqrt{f_0(t)} = f_0(t)/|\mu_0(t)| \leq 3f_0(t)/|\mu_0(0)| \leq 4f_0(t)n^{-3+\epsilon}$ by the first line of (4.47), we have for $t \geq t_0$,

$$\dot{f}_0(t) \geq 2^{-5-2N} \gamma_0 N^{-1} n^4 f_0 - \sqrt{f_0} C n^7 \geq 2^{-6-2N} \gamma_0 N^{-1} n^4 f_0,$$

if n is sufficiently small. Hence

$$f_0(t) \geq f_0(t_0) \exp \{2^{-6-2N} \gamma_0 N^{-1} n^4 (t - t_0)\}, \quad (4.62)$$

for $t \in [t_0, t'_1]$. However, using the definition of t'_1 and

$$f_0(t_0) \geq \frac{3}{4} |x_0(t_0)|^2 \geq \frac{3}{16} |x_0(0)|^2 \geq \frac{3}{16} n^{6-2\epsilon},$$

(4.62) implies $f_0(t'_1) > 2^{-2-N}n^2$, which is a contradiction. This contradiction shows the existence of $t_1 \in [t_0, t'_1]$ so that $f_0(t_1) = 2^{-2-N}n^2$. We also have (4.44) and (4.62) for all $t \leq t_1$.

Step 3. Transition regime (ii).

For $t \geq t_1$, $f_0(t)$ is large enough to cause $g(t)$ to decay in a rate we can control. Define

$$t_2 \equiv \inf_{t \geq t_1} \{t : g(t) \leq n^{2+2\sigma}\}. \quad (4.63)$$

We want to show that

$$t_1 \leq t_2 \leq t'_2 \equiv t_1 + n^{-4-2\sigma}. \quad (4.64)$$

Suppose the contrary, then $g(t) \geq n^{2+2\sigma}$ for all $t \leq t'_2$. By (4.47), for $t > t_1$ we have $\sqrt{f_0(t)} \leq f_0(t)(\frac{1}{2}f_0(t_1))^{-1/2} \leq 2^{(3+N)/2}nf_0(t)$. Recall from (4.61) that $\sum_{a,b=0}^N 2(\operatorname{Re} d_{ab}^0)f_a f_b \geq 2\gamma_0 N^{-1}g^2$. Hence for $t \in [t_1, t'_2]$,

$$\dot{f}_0(t) \geq 2\gamma_0 N^{-1}g^2 - \sqrt{f_0}Cn^7 > 0.$$

Thus $\dot{f}_0 > 0$ and $f_0(t) \geq 2^{-2-N}n^2$ for $t \in [t_1, t'_2]$. In particular (4.44) holds for $t > t_1$ and (4.45) holds for $x_0(t_2)$ should t_2 exist. By eq.(4.27) for $\dot{f}_0$,

$$\begin{aligned} \dot{g}(t) &\leq -\dot{f}_0(t) + \sum_{k=0}^N 2 \operatorname{Re} \bar{\mu}_k g_k \\ &= -\left\{ \sum_{a,b=0}^N 2(\operatorname{Re} d_{ab}^0)f_a f_b \right\} f_0 + \sum_{k=1}^N 2 \operatorname{Re} \bar{\mu}_k g_k \\ &\leq -2\gamma_0 N^{-1}g^2 f_0 + \sum_{k=1}^N 2 \operatorname{Re} \bar{\mu}_k g_k \\ &\leq -2^{-1-N}n^2\gamma_0 N^{-1}g^2 + Cn n^7 \\ &< -2^{-2-N}\gamma_0 N^{-1}n^2g^2, \end{aligned}$$

if n is sufficiently small. In the second line we have cancelled $2 \operatorname{Re} \bar{\mu}_0 g_0$. In the third line we used (4.61). Thus, by comparison principle,

$$g(t) < [g(t_1)^{-1} + 2^{-2-N}\gamma_0 N^{-1}n^2(t-t_1)]^{-1}, \quad (t_1 < t \leq t'_2).$$

for $t \in (t_1, t'_2]$. However, by the definition of t'_2 this implies $g(t'_2) < n^{2+2\sigma}$, which is a contradiction. This contradiction shows the existence of t_2 satisfying (4.64). Similarly, since $\dot{g}(t) > -Cn^2g(t)^2$ for $t_1 \leq t \leq t_2$, we have a lower bound $g(t) \geq [g(t_1)^{-1} + Cn^2(t-t_2)]^{-1}$ by comparison principle. This implies $t_2 - t_1 \geq Cn^{-4}$. We have proven (4.43)–(4.45) in Proposition 4.3 using assumption (4.47). Since (4.47) holds for $t = 0$, it holds for all $t \leq t_2$ by continuity. Hence (4.43)–(4.45) are proven.

Finally we prove (4.46) for $e^{-i\tau H_0}\xi(t_2)$. Let $t = t_2 + \tau$. We have

$$e^{-i\tau H_0}\xi(t_2) = e^{-itH_0}\xi_0 + J(t), \quad J(t) \equiv \int_0^{t_2} e^{-i(t-s)H_0} \mathbf{P}_c G(s) ds.$$

The estimate of $e^{-itH_0}\xi_0$ is by (4.42), since $\Lambda_5(t) \leq Cn^3(t_* + t_2)(1+t)^{-9/10}$ and $\Lambda(t) \leq Cn^3(t_* + t_2)(1+t)^{-1}(1+\tau)^{-1/2}$ if $t_2 \geq n^{-2}$. To estimate $J(t)$, we use the following integral inequalities: For $t \geq T \geq 1$,

$$\int_0^T |t-s|^{-9/10} ds \leq CTt^{-9/10}. \quad (4.65)$$

$$\int_0^T \min \{ (t-s)^{-3/2}, (t-s)^{-9/10} \} ds \leq CTt^{-1} \langle t-T \rangle^{-1/2}. \quad (4.66)$$

See [22, Lemma 2.6]. Since $\|G(s)\|_{L^1 \cap L^{5/4}} \leq Cn^3$ for $s \in [0, t_2]$, we have

$$\begin{aligned} \|J(t)\|_{L^5} &\leq C \int_0^{t_2} |t-s|^{-9/10} \|G_\xi(s)\|_{L^{5/4}} ds \leq C \int_0^{t_2} |t-s|^{-9/10} n^3 ds \\ &\leq Cn^3 t_2 (1+t)^{-9/10}. \end{aligned}$$

$$\begin{aligned} \|J(t)\|_{L^2_{\text{loc}}} &\leq C \int_0^{t_2} \min \{ |t-s|^{-3/2}, |t-s|^{-9/10} \} \|G(s)\|_{L^1 \cap L^{5/4}} ds \\ &\leq C \int_0^{t_2} \min \{ |t-s|^{-3/2}, |t-s|^{-9/10} \} n^3 ds \\ &\leq Cn^3 t_2 t^{-1} (1+t-t_2)^{-1/2}. \end{aligned}$$

The proof of Proposition 4.3 is complete. **Q.E.D.**

PROOF OF THEOREM 1.1 To prove Theorem 1.1 using Theorem 3.1, we want to use $\psi(t_2)$ as initial data and decompose $\psi(t_2)$ according to (3.1), i.e., with respect to $\mathcal{L}$, and check the assumptions (3.4)–(3.5). If the last condition in (4.41) fails, we simply set $t_2 = 0$. Let $\rho_0 = (|x_1(t_2)|^2 + \cdots + |x_N(t_2)|^2)^{1/2}$. We have

$$|x_0(t_2)| \sim n, \quad \rho_0 = (1 + o(1))n^{1+\sigma}, \quad \|\xi(t_2)\|_{L^2_{\text{loc}}} \leq Cn^3. \quad (4.67)$$

We may redefine $n = |x_0(t_2)|$. Let Q_{E_0} be the unique nonlinear ground state so that $Q_{E_0} = n\phi_0 + k$ with $k \perp \phi_0$, $k = O(n^3)$. Also choose $e^{i\theta_0} = n^{-1}x_0(t_2)$. We have $\|\psi(t_2) - Q_{E_0}e^{i\theta_0}\|_{L^2_{\text{loc}}} \leq \|k\|_{L^2_{\text{loc}}} + \|\psi(t_2) - x_0(t_2)\phi_0\|_{L^2_{\text{loc}}} \lesssim n^3 + \rho_0 + n^3 \lesssim \rho_0$, which is much smaller than n . By Lemma 2.2, for each E close to E_0 , we can decompose $\psi(t_2)$ as $\psi(t_2) = [Q_E + aR_E + \zeta + \eta]e^{i\theta}$ using the decomposition (3.1) with respect to $\mathcal{L}_E$. We have

$$e^{-i\theta}\psi(t_2) - Q_E = aR + \zeta + \eta = k + Z + \xi, \quad (4.68)$$

where $Z = e^{-i\theta}[x_1(t_2)\phi_1 + \cdots + x_N(t_2)]$ and $\xi = e^{-i\theta}\xi(t_2)$. With $E = E_0$ in (4.68), we have $a_{E_0} = (c_1Q, \text{Re } k + Z + \xi) = c_1(n\phi_0 + k, \text{Re } k + Z + \xi)$. Since $\phi_0 \perp (k + Z + \xi)$, we have

$$|a_{E_0}| \leq n^3(n^3 + \rho_0 + n^3) \lesssim Cn^3\rho_0,$$

which is much smaller than ρ_0^2 . Denote $P_{\mathbf{E}} = \sum_{k=1}^N P_{\mathbf{E}_k}$. We also have

$$\|\zeta - P_{\mathbf{E}}Z\| = \|P_{\mathbf{E}}(k + \xi)\| \lesssim n^3 + n^2 \|\xi\|_{L_{\text{loc}}^2} \lesssim n^3,$$

and hence $\|\zeta\| = \rho_0 + O(n^3)$. We have verified (3.4) with $E = E_0$. For all E with $|E - E_0| \leq D\rho_0^2$, write $\eta_1 = \mathbf{P}_c^{\mathcal{L}}(k + Z)$ and $\eta_2 = \mathbf{P}_c^{\mathcal{L}}\xi = \mathbf{P}_c^{\mathcal{L}}e^{-i\theta}\xi(t_2)$. We have

$$e^{s\mathcal{L}}\eta = e^{s\mathcal{L}}\eta_1 + e^{s\mathcal{L}}\eta_2.$$

Since η_1 is a local function bounded by $\|k\| + n^2\|Z\| \lesssim n^3$, we get

$$\begin{aligned} \|e^{s\mathcal{L}}\eta_1\|_{L^5} &\leq Cn^3(1+s)^{-9/10} \ll n^{4/5}\rho(s)^{8/5}, \\ \|e^{s\mathcal{L}}\eta_1\|_{L_{\text{loc}}^2} &\leq Cn^3(1+s)^{-3/2} \ll n^{1/2}(1+s)^{-1/2}\rho^2(s). \end{aligned}$$

Recall $\rho(s) = [\rho_0^{-2} + N^{-1}\gamma_0 n^2 s]^{-1/2}$ with $\rho_0 \sim n^{1+\sigma}$. Write $\mathcal{L}$ as $\mathcal{L} = -i(H_0 - E) + W$, where $Wh = 2\lambda Q^2 h + \lambda Q^2 \bar{h}$ is a local operator bounded by n^2 . We have

$$e^{t\mathcal{L}}\eta_2 = \mathbf{P}_c^{\mathcal{L}}e^{-it(H_0-E)}e^{-i\theta}\xi(t_2) + \int_0^t e^{(t-s)\mathcal{L}}We^{-is(H_0-E)}e^{-i\theta}\xi(t_2)ds.$$

Hence

$$\begin{aligned} \|e^{t\mathcal{L}}\eta_2\|_{L^5} &\leq \|e^{-itH_0}\xi(t_2)\|_{L^5} + \int_0^t |t-s|^{-9/10}n^2 \|e^{-isH_0}\xi(t_2)\|_{L_{\text{loc}}^2} ds, \\ \|e^{t\mathcal{L}}\eta_2\|_{L_{\text{loc}}^2} &\leq \|e^{-itH_0}\xi(t_2)\|_{L_{\text{loc}}^2} \\ &\quad + \int_0^t \min\{|t-s|^{-3/2}, |t-s|^{-9/10}\} n^2 \|e^{-isH_0}\xi(t_2)\|_{L_{\text{loc}}^2} ds. \end{aligned}$$

Using the out-going estimates (4.46) for $e^{-isH_0}\xi(t_2)$, we conclude the estimates (3.5) for η_E for all E close to E_0 with $|E - E_0| \leq D\rho_0^2$. **Q.E.D.**

Acknowledgments

The author would like to thank H.-T. Yau for his very helpful comments and discussions. The author was partially supported by NSF grant DMS-9729992.

References

- [1] V.S. Buslaev and G.S. Perel'man, Scattering for the nonlinear Schrödinger equations, states close to a soliton, St. Petersburg Math J. **4** (1993) 1111–1142.
- [2] V.S. Buslaev and G.S. Perel'man, On the stability of solitary waves for nonlinear Schrödinger equations. Nonlinear evolution equations, 75–98, Amer. Math. Soc. Transl. Ser. 2, 164, Amer. Math. Soc., Providence, RI, 1995.

- [3] V.S. Buslaev and C. Sulem, On the asymptotic stability of solitary waves of nonlinear Schrödinger equations, preprint.
- [4] S. Cuccagna, Stabilization of solutions to nonlinear Schrödinger equations. *Comm. Pure Appl. Math.* **54** (2001), no. 9, 1110–1145.
- [5] S. Cuccagna, On asymptotic stability of ground states of NLS, preprint.
- [6] M. Grillakis: Linearized instability for nonlinear Schrödinger and Klein-Gordon equations. *Comm. Pure Appl. Math.* **41** (1988), no. 6, 747–774.
- [7] M. Grillakis: Analysis of the linearization around a critical point of an infinite dimensional Hamiltonian system, *Comm. Pure Appl. Math.* **43** (1990), 299–333.
- [8] A. Jensen and T. Kato, Spectral properties of Schrödinger operators and time-decay of the wave functions, *Duke Math. J.* **46** no. 3, (1979), 583–611.
- [9] J.-L. Journe, A. Soffer and C.D. Sogge, Decay estimates for Schrödinger operators. *Comm. Pure Appl. Math.* **44** (1991), no. 5, 573–604.
- [10] C.-A. Pillet and C.E. Wayne, Invariant Manifolds for a class of dispersive, Hamiltonian, partial differential equations, *J. Diff. Equations* **141**, (1997), 310–326.
- [11] Rauch, J. Local decay of scattering solutions to Schrödinger’s equation. *Comm. Math. Phys.* **61** (1978), no. 2, 149–168.
- [12] H.A. Rose and M.I. Weinstein, On the bound states of the nonlinear Schrödinger equation with a linear potential, *Physica D* **30** (1988), 207–218.
- [13] J. Shatah and W. Strauss, Instability of nonlinear bound states. *Comm. Math. Phys.* **100** (1985), no. 2, 173–190.
- [14] J. Shatah and W. Strauss, Spectral condition for instability. *Nonlinear PDE’s, dynamics and continuum physics* (South Hadley, MA, 1998), 189–198, *Contemp. Math.*, 255, Amer. Math. Soc., Providence, RI, 2000.
- [15] I. M. Sigal, Non-Linear wave and Schrödinger equations I, instability of periodic and quasiperiodic solutions, *Comm. Math. Phys.* **153** (1993) 297–320.
- [16] A. Soffer and M.I. Weinstein, Multichannel nonlinear scattering theory for non-integrable equations I, II, *Comm. Math. Phys.* **133** (1990), 119–146; *J. Diff. Equations* **98**, (1992), 376–390.

- [17] A. Soffer and M.I. Weinstein, Resonances, radiation damping and instability in Hamiltonian nonlinear wave equations, *Invent. math.* **136**, (1999), 9–74.
- [18] A. Soffer and M.I. Weinstein, Selection of the ground state for nonlinear Schrödinger equations, preprint.
- [19] T.-P. Tsai and H.-T. Yau, Asymptotic dynamics of nonlinear Schrödinger equations, resonance dominated and dispersion dominated solutions, *Comm. Pure Appl. Math.* **55** (2002) 153–216.
- [20] T.-P. Tsai and H.-T. Yau, Relaxation of excited states in nonlinear Schrödinger equations, *IMRN*, to appear.
- [21] T.-P. Tsai and H.-T. Yau, Stable directions for excited states of nonlinear Schrödinger equations, *Comm. P.D.E.*, to appear.
- [22] T.-P. Tsai and H.-T. Yau, Classification of asymptotic profiles for nonlinear Schrödinger equations with small initial data, preprint.
- [23] M.I. Weinstein, Lyapunov stability of ground states of nonlinear dispersive evolution equations, *Comm. Pure Appl. Math.* **39**, (1986), 51–68.
- [24] K.Yajima, The $W^{k,p}$ continuity of wave operators for Schrödinger operators, *J.Math. Soc. Japan* **47**, no. 3, (1995), 551–581.